

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
10 May 2002 (10.05.2002)

PCT

(10) International Publication Number  
**WO 02/36824 A1**

(51) International Patent Classification<sup>7</sup>: **C12Q 1/68**,  
C12N 15/12, A01K 67/027, C07K 16/40

(74) Agents: CALHOUN, Douglas, C. et al.; A J Park, Hud-  
dard Parker Building, 6th Floor, Post Office Square, P.O.  
Box 949, 6015 Wellington (NZ).

(21) International Application Number: PCT/NZ01/00245

(22) International Filing Date: 31 October 2001 (31.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
507888 31 October 2000 (31.10.2000) NZ  
508662 6 December 2000 (06.12.2000) NZ

(71) Applicants and

(72) Inventors: **GEORGES, Michel, Alphonse, Julien**  
[BE/BE]; 24 rue Vieux Tige, B-4161 Villers-aux-Tours  
(BE). **COPPIETERS, Wouter, Herman, Robert**  
[BE/BE]; Beekstraat 2, B-3401 Landen (BE). **GRISART,**  
**Bernard, Marie-Josée, Jean** [BE/BE]; 2 Rue Bordia,  
B-4218 Couthuin (BE). **SNELL, Russell, Grant** [NZ/NZ];  
3/28 Tennyson Street, Balmoral, Auckland (NZ). **REID,**  
**Suzanne, Jean** [NZ/NZ]; ViaLactia Biosciences (NZ)  
Limited, P O Box 109-185, Newmarket, Auckland (NZ).  
**FORD, Christine, Ann** [NZ/NZ]; ViaLactia Biosciences  
(NZ) Limited, P O Box 109/185, Newmarket, Auckland  
(NZ). **SPELMAN, Richard, John** [NZ/NZ]; 340 Horotiu  
Road, R.D. 8, Hamilton (NZ).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,  
YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



WO 02/36824 A1

(54) Title: MARKER ASSISTED SELECTION OF BOVINE FOR IMPROVED MILK PRODUCTION USING DIACYLGLYC-  
EROL ACYLTRANSFERASE GENE DGAT1

(57) Abstract: The present invention provides a method of genotyping bovine for improved milk production traits by determining the DGAT1 genotypic state of said bovine, wherein the DGAT1 gene and polymorphisms within said gene have been found to be associated with such improved milk production traits.

**Marker Assisted Selection of Bovine for Improved Milk Production Using Diacylglycerol  
Acyltransferase gene DGAT1**

**FIELD OF THE INVENTION**

5 This invention relates to an application of marker assisted selection of bovine for a quantitative trait loci (QTL) associated with milk production, particularly although by no means exclusively, by assaying for the presence of at least one allele which is associated with increased milk volume as well as improved milk composition. The present invention also relates to the gene associated with the QTL, various  
10 polymorphisms within the gene sequence, proteins encoded by these sequences as well as to the application of all of these in the farming industry.

**BACKGROUND**

15 The genetic basis of bovine milk production is of immense significance to the dairy industry. An ability to modulate milk volumes and content has the potential to alter farming practices and to produce products which are tailored to meet a range of requirements. In particular, a method of genetically evaluating bovine to select those which express desirable traits, such as increased milk production and improved milk  
20 composition, would be desirable.

To date, bovine genomics are poorly understood and little is known regarding the genes which are critical to milk production. While there have been reports of quantitative trait loci (QTLs) on bovine chromosome 14 postulated to be associated with milk production  
25 (Coppieters *et al* (1998)), the specific genes involved have not to date been identified.

Marker assisted selection, which provides the ability to follow a specific favourable genetic allele, involves the identification of a DNA molecular marker or markers that segregate with a gene or group of genes associated with a QTL. DNA markers have  
30 several advantages. They are relatively easy to measure and are unambiguous, and as DNA markers are co-dominant, heterozygous and homozygous animals can be distinctively identified. Once a marker system is established, selection decisions are able to be made very easily as DNA markers can be assayed at any time after a DNA containing sample has been collected from an individual infant or adult animal, or even  
35 earlier as it is possible to test embryos in vitro if such embryos are collected.

The applicants have now identified a gene responsible for the QTL effect on bovine chromosome 14 as well as a number of polymorphisms which are associated with distinct genetic merits of animals for milk composition and volume.

- 5 It is an object of the present invention to provide an application method for marker assisted selection of this bovine gene, and in particular, of the polymorphisms in the bovine gene which are associated with increased milk volume and altered milk composition; and/or to provide genetic markers for use in such a method; and/or to provide the nucleic acid and amino acid sequences of this gene and encoded  
10 polypeptide; and/or to provide animals selected using the method of the invention as well as milk produced by the selected animals; and/or to provide the public with a useful choice.

#### **SUMMARY OF THE INVENTION**

- 15 This invention relates to the discovery of the bovine Diacylglycerol-o-acyltransferase (DGAT1) gene and polymorphisms within the bovine DGAT1 gene which are associated with increased milk yield and altered milk composition.
- 20 More specifically, several polymorphisms in the bovine DGAT1 gene have been identified distinguishing multiple DGAT1 alleles in different cattle breeds. These polymorphisms include: K232A (Bases 6829/30 AA-CG nucleic acid change and K-A amino acid change); Nt984+8(Base 7438 A-G nucleic acid change); Nt984+26(Base 7456 C-T nucleic acid change); Nt1470+85(Base 8402 C-T nucleic acid change); Nt191+435 (Base  
25 626 T-G nucleic acid change); Nt191-3321 (Base 3512 T-G nucleic acid change); Nt279+144 (Base 4040 T-C nucleic acid change); Nt279+1067 (Base 4963 A-G nucleic acid change); Nt279+1107 (Base 5003 G-A nucleic acid change); Nt358 (Base 5997 C-T nucleic acid change); Nt754+3 (Base 6892 G-A nucleic acid change); Nt897+32 (Base 7224/5 GG-AC nucleic acid change); Nt1251+42 (Base 7987 G-A nucleic acid change)  
30 as summarised in Table 1. In particular, DGAT1 alleles characterized by the K232A mutation have been identified as being associated with an increased milk volume and altered milk composition in animals dependent upon whether they are homozygous with or without the mutation or heterozygous carrying one mutated allele. More specifically, the presence of the K232A mutation results in a decrease in milkfat percentage, milkfat  
35 yield, solid fat content and milk protein percentage, while increasing milk volume and milk protein yield.

The present invention thus relates to the use of the polymorphisms in a method of identification and selection of a bovine having at least one of said polymorphisms as well as to providing markers specific for such identification. Kits comprising said markers for use in marker selection also form part of the present invention as do  
5 animals so selected, the milk produced by such selected animals and products produced from such milk, particularly as such milk and milk products affect processing and/or health characteristics for consumers.

In particular, the present invention is directed to a method of genotyping cows or bulls  
10 for one or more of the polymorphisms disclosed herein, selected cows or bulls so genotyped and milk and semen from said selected cows and bulls respectively.

According to a further aspect the present invention is directed to the isolated DGAT1 nucleic acid and allelic nucleic acid molecules comprising polymorphisms as well as to  
15 the proteins encoded thereby and their polypeptide sequences. Antibodies raised against said proteins are also contemplated, as are vectors comprising the nucleic acid molecules, host cells comprising the vectors; and protein molecules expressed in said host cells; and the application of all of them in the farming industry.

20 In particular, such applications include methods for modulating milk production and/or composition in a lactating bovine by affecting DGAT1 activity, by reducing the activity of DGAT1 (e.g. by use of specific ribozymes, antisense sequences and/or antibodies, or by transgenic technology to produce a "knock out" bovine and/or bovine with introduced transgenes containing the DGAT1 gene and/or variations of this gene  
25 driven by various promoters).

#### BRIEF DESCRIPTION OF THE DRAWINGS

The invention will now be described with reference to the Figures of the accompanying  
30 drawings in which:

**Figure1:** Shows a BAC contig spanning the *BULGE13-BULGE09* interval relative to a schematic diagram of bovine chromosome 14 and a schematic diagram showing the location of the genetic markers. The most likely position of the QTL is shown as a bar  
35 on the FISH-anchored linkage map proximal to BTA14q. The BACs composing the contigs spanning the *BULGE13-BULGE09* interval are shown as a series of horizontal lines. The symbols on each BAC indicate their individual STS content: solid circles

correspond to STS derived from BAC ends, open boxes to microsatellite markers, and solid triangles to gene-specific Comparative Anchored Tagged Sequences. The arrow heads mark the BACs from which the respective BAC end STS were derived. The length of the lines do not reflect the actual insert size of the corresponding BACs. The BAC

5 contig was aligned with the orthologous human HSA8q24.3 genomic "golden path" sequence represented according to the Ensembl Human Genome Server (<http://www.ensembl.org/>): individual sequence contigs are shown in alternating light and dark; a horizontal line indicates a gap in the sequence assembly; genetic markers are indicated under the contig map; the lines and boxes above the contig map represent

10 "curated", "predicted known" or "predicted novel" genes.

**Figures 2a and 2b:** Show the genomic sequence of the bovine DGAT1 gene. Figure 2a is the 31 base pair sequence upstream but adjacent to the ATG or translation start site and is 5'UTR. Figure 2b is the genomic sequence in the bovine DGAT1 gene from the

15 ATG translation start site (base 1) through to genomic sequence flanking the gene at the 3' end. The significant features including intron/exon boundaries, polymorphic sites, polyadenylation signal, and alternate splicing site and some of the primer sequences used in the assays described herein, are indicated;

20 **Figure 3:** Shows the genomic organization, four polymorphisms and haplotypes found in the bovine *DGAT1* gene. Leader and trailer sequences are shown in light grey, coding sequences in dark grey and intronic sequences as a hollow line. The positions of four of the identified polymorphisms are marked as shown on the gene, and detailed in the underlying boxes including the corresponding sequence traces All the sequence

25 variations are summarised in Table 1. The four *DGAT1* haplotypes which were found in the Dutch and New-Zealand Holstein-Friesian population as defined by these polymorphisms are shown and referred to as "*sHQ-D*", "*sHQ-NZ*", "*sHQ-III*" for the fat increasing haplotypes and "*shq*" for the fat decreasing haplotype;

30 **Figure 4a:** Shows the corresponding full length amino acid sequence for DGAT1 sequence of Figure 2b including annotation of the amino acid substitution;

**Figure 4b:** Shows the amino acid sequence predicted as a result of alternate splicing with exon VIII;

35

**Figure 5:** Shows the multiple peptide alignment of a portion of the *DGAT1* protein flanking the K232A substitution from *Bos taurus*, *Bison bison*, *Ovis aries*, *Sus scrofa*,

*Homo sapiens*, *Cercopithecus aethiops*, *Mus musculus domesticus* and *Rattus norvegicus* showing the evolutionary conservation of the lysine mutated in the bovine K232A polymorphism;

- 5 **Figure 6:** A. Shows the frequency distribution of observed *DGAT1* SNP haplotypes in the Dutch and New Zealand Holstein-Friesian dairy cattle populations. B-D. Shows the frequency distribution of the combined microsatellite (*BULGE09-BULGE11*) and SNP *DGAT1* haplotypes. The *HQ-D* and *HQ-NZ* haplotypes are shown; and
- 10 **Figure 7:** Shows the lod score due to LD when including (+) or excluding (-) the four *DGAT1* polymorphisms shown in Figure 3 in a combined linkage and LD multipoint maximum likelihood mapping method. The lod score corresponds to the  $\log_{10}$  of the ratio between the likelihood of the data assuming LD and linkage between the markers and the likelihood of the data assuming linkage in the absence of LD. The positions of
- 15 the microsatellites and SNP markers utilized in the analysis are shown on the X-axis, while the position of the *DGAT1* SNPs is marked by a red arrow at the top of the figure.

#### DETAILED DESCRIPTION OF THE INVENTION

- 20 It has been discovered for the first time that the *DGAT1* gene in bovine is associated with the QTL on chromosome 14 which is linked with improved milk production traits. More particularly, a number of novel polymorphisms on the *DGAT1* gene have been discovered. It is thought that one or more of these polymorphisms is responsible for these traits.

25

- The method used for isolating genes which cause specific phenotypes is known as positional candidate cloning. It involves: (i) the chromosomal localisation of the gene which causes the specific phenotype using genetic markers in a linkage analysis; and (ii) the identification of the gene which causes the specific phenotype amongst the
- 30 "candidate" genes known to be located in the corresponding region. Most of the time these candidate genes are selected from available mapping information in humans and mice.

- The tools required to perform the initial localisation (step (i) above) are microsatellite marker maps, which are available for livestock species and are found in the public domain (Bishop *et al.*, 1994; Barendse *et al.*, 1994; Georges *et al.*, 1995; and Kappes, 1997). The tools required for the positional candidate cloning, particularly the BAC libraries, (step (ii) above) are partially available from the public domain. Genomic
- 35

libraries with large inserts constructed with Bacterial Artificial Chromosomes (BAC) are available in the public domain for most livestock species including cattle. For general principles of positional candidate cloning, see Collins, 1995 and Georges and Anderson, 1996.

5 Recently, a quantitative trait locus (QTL) with major effect on milk solids composition, located at the centromeric end of bovine chromosome 14, has been reported (Coppieters *et al.*, (1998)). This QTL was shown to effect milk fat content and in particular to significantly affect protein %, volume, protein yield and fat yield of milk. The linkage  
10 study as well as subsequent marker assisted segregation analyses allowed for the identification of thirteen Holstein-Friesian sires predicted to be heterozygous "Qq" for the corresponding QTL (Coppieters *et al.*, (1998); Riquet *et al.*, (1999)).

Linkage disequilibrium methods were applied to refine the map position of the QTL to a  
15  $\approx 5$  cM interval bounded by microsatellite markers BULGE09 and BULGE30.

A bovine DGAT1 nucleotide sequence was determined by the applicants and is shown in Figures 2a and 2b with the corresponding amino acid sequences (long and short forms) being shown in Figures 4a and 4b respectively. Table 1 sets out all the polymorphisms  
20 located to date with reference to the sequence in Figure 2b. Some of the genetic polymorphisms identified in the bovine DGAT1 gene are reported in Figure 3. The nucleic acid and protein sequences of the DGAT1 alleles including the K232A mutation are shown in Figures 2a and 2b (SEQ ID NOs: 3 and 1), annotated to show the alternatively spliced forms. The cDNA sequence is also set out in SEQ ID NO: 4.

25 The sequence information in the Figures gives rise to numerous, and separate, aspects of the invention.

In one aspect, the invention provides a method of determining genetic merit of a bovine  
30 with respect to milk composition and volume which comprises the step of determining the bovine DGAT1 genotypic state of said bovine. In particular, this method is useful for genotyping and selecting cows and bulls having the desired genotypic state so that milk and semen may be collected from said cows and bulls respectively. Such semen would be useful for breeding purposes to produce bovine having the desired genotypic  
35 and, as a result, phenotypic state. In addition, cows genotyped by the methods of the present invention are also useful for breeding purposes, particularly for breeding with the selected bulls and/or to be artificially inseminated with the semen from selected

bulls. The embryos and offspring produced by such cows also form part of the present invention.

In one embodiment, the genotypic state is determined with respect to DNA obtained  
5 from said bovine.

Alternatively, said genotypic state is determined with reference to mRNA obtained from said bovine.

10 In yet a further embodiment, the genotypic state is determined with reference to the amino acid sequence of expressed bovine DGAT1 protein obtained from said bovine.

Conveniently, in said method, the genotypic state of DNA encoding bovine DGAT1 is determined, directly or indirectly.

15

Alternatively, in said method the genotypic state of at least one nucleotide difference from the nucleotide sequence encoding bovine DGAT1 is determined, directly or indirectly.

20 More specifically, in said method the genotypic state of bovine DGAT1 allele(s) characterised by one or more of the polymorphisms shown in Table 1 below, is determined, directly or indirectly.



**Table 1:**  
**Table of polymorphisms in the bovine DGAT1 gene**

5 Start codon (atg); the a residue is denoted as position 1

Base number relative to exonic sequence <sup>1</sup>	Nucleotide distance from start	substitution	Intron/exon #
Nt 191 + 435	626	T-G CAGTGCTAGGGG CAGTGCGAGGGG	Intron 1
Nt 191 + 3321	3512	T-G GCATTGCGCT GCATGGCGCT	Intron 1
Nt 279 + 144	4040	T-C TACCCCTGGGAC TACCCCGGGAC	Intron 2
Nt 279 + 1067	4963	A-G CTCTTAGCAGC CTCTTGGCAGC	Intron 2
Nt 279 + 1107	5003	G-A ACAGGCAACT ACAGACAACT	Intron 2
Nt 358	5997	C-T TGTCTCTGTTC TGTCTTTGTTC	Exon IV
Nt 692	6829	AA-GC GGTAAGAAGGCCAA (Q) GGTAAGGCGGCCAA (q)	K232A Exon VIII *
Nt 754 + 3	6892	G-A GCGGTGAGGAT GCGGTAAGGAT	Intron VIII
Nt 897 + 32	7224	GG-AC GGGGGGGGGGGACTCT GGGGGACGGGGACTCT	Intron X
Nt 984 + 8	7438	A-G GAGTGACCTGC GAGTGGCCTGC	Intron XII *
Nt 984 + 26	7456	C-T GGACGCGTGGG GGACGTGTGGG	Intron XII *
Nt1251 + 42	7987	G-A GGTGGGGGTGG GGTGGAGGTGG	Intron XV
Nt 1470 + 85	8402	C-T CTGGGCGCAGC CTGGGTGCAGC	3' flanking region *

The numbers given are for the actual nucleotide or in the case of two nucleotide substitutions to the first nucleotide in the variation (counting 5' to 3')

10 \*More detail of these polymorphisms is given in Figure 2b.

<sup>1</sup> e.g. Nt 191 represents nucleotide number 191 from the start site of the coding sequence, + 435 represents number of nucleotides from and including base 192 in the genomic sequence (intron 1) to the polymorphic nucleotide. The polymorphic nucleotides are shaded

15 Preferably, the invention is directed to a method of determining the genotypic state of bovine DGAT1 allele(s) by determining the presence of the K232A polymorphism, either directly or indirectly.

There are numerous art standard methods known for determining whether a particular  
20 DNA sequence is present in a sample. An example is the Polymerase Chain Reaction

(PCR). A preferred aspect of the invention thus includes a step in which ascertaining whether a polymorphism(s) in the sequence of DGAT1 DNA is present, includes amplifying the DNA in the presence of primers based on the nucleotide sequence of the DGAT1 gene and flanking sequence, and/or in the presence of a primer containing at least a portion of a polymorphism as disclosed herein and which when present results in altered relative milk lipid and protein production, and milk volume.

A primer of the present invention, used in PCR for example, is a nucleic acid molecule sufficiently complementary to the sequence on which it is based and of sufficient length to selectively hybridise to the corresponding portion of a nucleic acid molecule intended to be amplified and to prime synthesis thereof under *in vitro* conditions commonly used in PCR. Likewise, a probe of the present invention, is a molecule, for example a nucleic acid molecule of sufficient length and sufficiently complementary to the nucleic acid molecule of interest, which selectively binds under high or low stringency conditions with the nucleic acid sequence of interest for detection thereof in the presence of nucleic acid molecules having differing sequences.

In another aspect, the invention provides a method for determining the genetic merit of bovine with respect to milk content and volume with reference to a sample of material containing mRNA obtained from the bovine. This method includes ascertaining whether a polymorphism(s) in the sequence of the mRNA encoding DGAT1 is present. The presence of such polymorphisms again indicates an association with altered relative milk lipid and protein production and milk volume.

Again, if an amplification method such as PCR is used in ascertaining whether a polymorphism(s) in the sequence of the mRNA encoding (DGAT1) is present, the method includes reverse transcribing the mRNA using a reverse transcriptase to generate a cDNA and then amplifying the cDNA in the presence of a pair of primers complementary to a nucleotide sequence encoding a protein having biological activity of wild type DGAT1.

In a further aspect, the invention includes the use of a probe in the methods of genotyping according to the invention wherein the probe is selected from any 5 or more contiguous nucleotides of the DGAT1 sequence as shown in Figure 2b, which is therefore sufficiently complementary with a nucleic acid sequence encoding such bovine DGAT1, or its complement, so as to bind thereto under stringent conditions. Diagnostic kits containing such a probe are also included. Such probes may be selected from

ForAA (FAM): CGTTGGCCTTCTTA or DgatADGC (VIC): TTGGCCGCCTTACC. (SEQ ID NOs: 20 and 21 respectively.)

5 The invention further includes isolated nucleic acid molecules encoding the DGAT1 variant proteins i.e. those proteins encoded by SEQ ID NOs: 1 and 4 (Figure 2b), with or without one or more polymorphisms of SEQ ID Nos: 7 to 19 (Table 1), or a fragment or variant thereof. Particularly, the invention includes an isolated nucleic acid molecule comprising a DNA molecule having in whole or in part the nucleotide sequence identified in Figure 2b or which varies from the sequence due to the degeneracy of the  
10 genetic code, or a nucleic acid strand capable of hybridising with said nucleic acid molecule under stringent hybridisation conditions.

The invention includes isolated mRNA transcribed from DNA having a sequence which corresponds to a nucleic acid molecule of the invention.

15

The invention includes isolated DNA in a recombinant cloning vector and a prokaryotic or eukaryotic cell containing and expressing heterologous DNA of the invention.

20 The invention includes a transfected cell line which expresses a protein encoded by the nucleic acid molecules of the invention.

The invention also includes a primer composition useful for detection of the presence of DNA encoding DGAT1 and/or the presence of DNA encoding a variant protein. In one form, the composition can include a nucleic acid primer substantially complementary to  
25 a nucleic acid sequence encoding DGAT1. The nucleic acid sequence can in whole or in part be that identified in Figure 2b. Diagnostic kits including such a composition are also included.

30 The invention further provides a diagnostic kit useful in detecting DNA encoding a variant DGAT1 protein in bovine which includes first and second primers for amplifying the DNA, the primers being complementary to nucleotide sequences of the DNA upstream and downstream, respectively, of a polymorphism in the portion of the DNA encoding DGAT1 which results in altered relative milk lipid, solid fat content and protein production and milk volume, wherein at least one of the nucleotide sequences is  
35 selected to be from a non-coding region of the DGAT1 gene. The kit can also include a third primer complementary to a polymorphism, disclosed herein, located on the DGAT1 gene.

The invention includes a process for producing a protein of the invention, including preparing a DNA fragment including a nucleotide sequence which encodes the protein; incorporating the DNA fragment into an expression vector to obtain a recombinant DNA molecule which includes the DNA fragment and is capable of undergoing replication; transforming a host cell with the recombinant DNA molecule to produce a transformant which can express the protein; culturing the transformant to produce the protein; and recovering the protein from resulting cultured mixture.

Thus in a further aspect, the invention provides a purified protein having biological activity of DGAT1. The terms "isolated" and "purified" as used herein, each refer to a protein substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesised. In certain preferred embodiments, the protein having biological activity of DGAT1 comprises an amino acid sequence and variants shown in Figures 4a and 4b (SEQ ID NOs: 2, 5 and 6). Furthermore, proteins having biological activity of DGAT1 that are encoded by nucleic acids which hybridise under stringent conditions to a nucleic acid comprising a nucleotide sequence shown in Figure 2b (SEQ ID NOs: 1 and 4) are encompassed by the invention.

Proteins of the invention having DGAT1 activity can be obtained by expression of a nucleic acid coding sequence in a suitable host cell using techniques known in the art. Suitable host cells include prokaryotic or eukaryotic organisms or cell lines, for example, yeast, *E. coli*, insect cells and COS1 cells. The recombinant expression vectors of the invention can be used to express a protein having DGAT1 activity in a host cell in order to isolate the protein. The invention provides a method of preparing a purified protein of the invention comprising introducing into a host cell a recombinant nucleic acid encoding the protein, allowing the protein to be expressed in the host cell and isolating and purifying the protein. Preferably, the recombinant nucleic acid is a recombinant expression vector. Proteins can be isolated from a host cell expressing the protein and purified according to standard procedures of the art, including ammonium sulfate precipitation, column chromatography (eg. ion exchange, gel filtration, affinity chromatography, etc.) electrophoresis, and ultimately, crystallisation (see generally "Enzyme Purification and Related Techniques". *Methods in Enzymology*, 22, 233-577 (1971)).

Alternatively, the protein or parts thereof can be prepared by chemical synthesis using techniques well known in the chemistry or proteins such as solid phase synthesis (Merrifield, 1964), or synthesis in homogeneous solution (Houbenwcy1, 1987).

5 It will of course be understood that a variety of substitutions of amino acids is possible while preserving the structure responsible for activity of the DGAT1 proteins disclosed herein. Conservative substitutions are described in the patent literature, as for example, in United States Patent No 5,264,558 or 5,487,983. It is thus expected, for example, that interchange among non-polar aliphatic neutral amino acids, glycine,  
10 alanine, proline, valine and isoleucine, would be possible. Likewise, substitutions among the polar aliphatic neutral amino acids, serine, threonine, methionine, asparagine and glutamine could possibly be made. Substitutions among the charged acidic amino acids, aspartic acid and glutamic acid, could probably be made, as could substitutions among the charged basic amino acids, lysine and arginine. Substitutions  
15 among the aromatic amino acids, including phenylalanine, histidine, tryptophan and tyrosine would also likely be possible. These sorts of substitutions and interchanges are well known to those skilled in the art. Other substitutions might well be possible. Of course, it would also be expected that the greater percentage of homology ie. sequence similarity, of a variant protein with a naturally occurring protein, the greater  
20 the retention of activity.

A further advantage may be obtained through chimeric forms of the proteins, as known in the art. A DNA sequence encoding each entire protein, or a portion of the protein, could be linked, for example, with a sequence coding for the C-terminal portion of *E. coli*  
25  $\beta$ -galactosidase to produce a fusion protein.

The proteins of the invention, or portions thereof, have numerous applications in turn. By way of example, each protein can be used to prepare antibodies which bind to a distinct epitope in an unconserved region of the protein. An unconserved region of the  
30 protein is one which does not have substantial sequence homology to other proteins.

Still further, the invention includes an antibody to a bovine DGAT1 variant protein encoded by a nucleotide sequence of the present invention as well as a diagnostic kit containing such an antibody.

35 Conventional methods can be used to prepare the antibodies. For example, by using a DGAT1 peptide, polyclonal antisera or monoclonal antibodies can be made using

standard methods. A mammal, (eg. a mouse, hamster, or rabbit) can be immunised with an immunogenic form of the peptide which elicits an antibody response in the mammal. Techniques for conferring immunogenicity on a peptide include conjugation to carriers or other techniques well known in the art. For example, the peptide can be administered in the presence of adjuvant. The progress of immunisation can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassay can be used to assess the levels of antibodies. Following immunisation, antisera can be obtained and, if desired, polyclonal antibodies isolated from the sera.

To produce monoclonal antibodies, antibody producing cells (lymphocytes) can be harvested from an immunised animal and fused with myeloma cells by standard somatic cell fusion procedures, thus immortalising these cells and yielding hybridoma cells. Such techniques are well known in the art. For example, the hybridoma technique originally developed by Kohler and Milstein (Kohler, 1975) as well as other techniques such as the human B-cell hybridoma technique (Kozbor, 1983) and screening of combinatorial antibody libraries (Huse, 1989). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with the peptide, and monoclonal antibodies isolated.

The term "antibody" as used herein is intended to include fragments thereof which are also specifically reactive with the target protein. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for whole antibodies. For example,  $F(ab)_2$  fragments can be generated by treating antibody with pepsin. The resulting  $F(ab)_2$  fragment can be treated to reduce disulfide bridges to produce Fab' fragments.

Another method of generating specific antibodies, or antibody fragments, reactive against the target proteins is to screen expression libraries encoding immunoglobulin genes, or portions thereof, expressed in bacteria, with peptides produced from the nucleic acid molecules of the present invention. For example, complete Fab fragments, VH regions and FV regions can be expressed in bacteria using phage expression libraries. See for example Ward *et al.*, Huse *et al.*, and McCafferty *et al.* (Ward, 1989); Huse 1989; McCafferty, 1990). Screening such libraries with, for example, a DGAT1 protein can identify immunoglobulin fragments reactive with that DGAT1. Alternatively, the SCID-hu mouse developed by Genpharm can be used to produce antibodies, or fragments thereof.

The polyclonal, monoclonal or chimeric monoclonal antibodies can be used to detect the proteins of the invention, portions thereof or closely related isoforms in various biological materials. For example, they can be used in an ELISA, radioimmunoassay or histochemical tests. Thus, the antibodies can be used to quantify the amount and location of a DGAT1 protein of the invention, portions thereof or closely related isoforms in a sample in order to determine the role of DGAT1 proteins. Using methods described hereinbefore, polyclonal, monoclonal antibodies, or chimeric monoclonal antibodies can be raised to non-conserved regions of DGAT1 and used to distinguish a particular DGAT1 from other proteins.

The polyclonal or monoclonal antibodies can be coupled to a detectable substance or reporter system. The term "coupled" is used to mean that the detectable substance is physically linked to the antibody. Suitable detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, and acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride and phycoerythrin; an example of a luminescent material includes luminol; and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$  and  $^3\text{H}$ . In a preferred embodiment, the reporter system allows quantitation of the amount of protein (antigen) present.

Such an antibody-linked reported system could be used in a method for determining whether a fluid or tissue sample of a bovine contains a deficient amount or an excessive amount of the relevant DGAT1 protein. Given a normal threshold concentration of such a protein, test kits can be developed.

The availability of such antibodies gives rise to further applications. One is a diagnostic kit for identifying cells comprising an antibody (such as a monoclonal antibody) which binds to a protein comprising an amino acid sequence shown in Figure 4a and 4b; means for detecting the antibody when bound to the protein, unreacted protein or unbound antibody; means for determining the amount of protein in the sample; and means for comparing the amount of protein in the sample with a standard. In some embodiments of the invention, the detectability of the antibody which binds to a specific DGAT1 protein is activated by the binding (eg. change in fluorescence spectrum, loss of

radioisotopic label). The diagnostic kit can also contain an instruction manual for use of the kit.

Antibody-based diagnostics are of course not the only possibility. A further diagnostic kit comprises a nucleotide probe complementary to the sequence, or an oligonucleotide fragment thereof, shown in Figure 2a and 2b, for example, for hybridisation with mRNA from a sample of cells; means for detecting the nucleotide probe bound to mRNA in the sample with a standard. In a particular aspect, the kit of this aspect of the invention includes a probe having a nucleic acid molecule sufficiently complementary with a sequence identified in Figure 2a and 2b, or its complement, so as to bind thereto under stringent conditions. "Stringent hybridisation conditions" takes on its common meaning to a person skilled in the art. Appropriate stringency conditions which promote nucleic acid hybridisation, for example, 6x sodium chloride/sodium citrate (SSC) at about 45°C are known to those skilled in the art, including in Current Protocols in Molecular Biology, John Wiley & Sons, NY (1989). Appropriate wash stringency depends on degree of homology and length of probe. If homology is 100%, a high temperature (65°C to 75°C) may be used. If homology is low, lower wash temperatures must be used. However, if the probe is very short (<100bp), lower temperatures must be used even with 100% homology. In general, one starts washing at low temperatures (37°C to 40°C), and raises the temperature by 3-5°C intervals until background is low enough not to be a major factor in autoradiography. The diagnostic kit can also contain an instruction manual for use of the kit.

One of the major applications of the present invention is in the marker assisted selection of bovines having a polymorphism in the DGAT1 gene and which are associated with improved milk production traits. The invention therefore provides a diagnostic kit which can be used to determine the DGAT1 genotype of bovine genetic material, for example. One kit includes a set of primers used for amplifying the genetic material. A kit can contain a primer including a nucleotide sequence for amplifying a region of the genetic material containing one of the polymorphisms described herein. Such a kit could also include a primer for amplifying the corresponding region of the normal DGAT1 gene, i.e. the sequence without polymorphisms. Usually, such a kit would also include another primer upstream or downstream of the region of interest complementary to a coding and/or non-coding portion of the gene. These primers are used to amplify the segment containing the mutation, i.e. polymorphism, of interest.



In particular, the invention is directed to the use of the polymorphisms in the DGAT1 gene in the genotyping of cows and bulls as well as to cows and bulls selected by such genotyping which have one or more of said polymorphisms in the DGAT1 gene. Such bulls so selected are of valuable breeding stock and the invention is also directed to the semen produced by such selected bulls for breeding purposes. Cows so selected are also useful as breeding stock as are their offspring. In addition, such cows may produce valuable dairy herds as the milk produced by such cows is produced in greater volumes than equivalent non-selected cows, and/or has an altered composition in that it comprises less milkfat and more milk protein. Such milk and products made therefrom also form part of the invention. It is also noted that the milk from these selected cows will be valuable as the fat content is not only decreased but is also characterised by being softer. Without being bound by theory, it is thought that this increased fat softness is due to the fatty acid composition being such that there is less saturated and more unsaturated fat in the milk of selected cows. Thus it is anticipated that products made from such milk will have processing advantages, such as in the production of more spreadable butter, as well as having a health benefit on consumers, as generally unsaturated fats are considered to be more "healthy" than saturated fats. The protein composition of milk produced by such selected cows is also altered. In particular, such milk comprises an altered protein yield compared to milk for non-selected cows and the casein:whey ratio is also altered which makes such milk valuable for cheese production.

Thus, the present invention involves genotyping bovine, both cows and bulls, for the DGAT1 polymorphisms disclosed herein, selected cows and bulls so genotyped, milk and semen produced by the selected cows and bulls so genotyped, offspring produced by the selected bovine, including embryos and cells (including cell lines) useful for cloning said selected bovine.

The actual genotyping is carried out using primers that target specific polymorphisms as described herein and that could function as allele-specific oligonucleotides in conventional hybridisation, Taqman assays, OLA assays, etc. Alternatively, primers can be designed to permit genotyping by microsequencing.

One kit of primers can include first, second and third primers, (a), (b) and (c), respectively. Primer (a) is based on a region containing a DGAT1 mutation such as described above. Primer (b) encodes a region upstream or downstream of the region to be amplified by primer (a) so that genetic material containing the mutation is amplified,

by PCR, for example, in the presence of the two primers. Primer (c) is based on the region corresponding to that on which primer (a) is based, but lacking the mutation. Thus, genetic material containing the non-mutated region will be amplified in the presence of primers (b) and (c). Genetic material homozygous for the DGAT1 gene will thus provide amplified products in the presence of primers (b) and (c). Genetic material  
5 homozygous for the mutated gene will thus provide amplified products in the presence of primers (a) and (b). Heterozygous genetic material will provide amplified products in both cases.

- 10 The present invention also contemplates the modulation of milk production and content in non-human animals by modulating the activity of the DGAT1 protein. In particular, this aspect of the invention includes a method of modulating milk production and/or milk content in a lactating bovine, the method comprising administering to the bovine an effective amount of a nucleic acid molecule substantially complementary to at least a  
15 portion of mRNA encoding the bovine DGAT1 variant proteins and being of sufficient length to sufficiently reduce expression of said DGAT1, i.e. by use of antisense nucleic acids.

- Antisense nucleic acids or oligonucleotides (RNA or preferably DNA) can be used to  
20 inhibit DGAT1 production in a bovine if this is considered desirable e.g. in order to produce a bovine capable of improved milk production, i.e. increased milk volume and decreased milkfat content. Antisense oligonucleotides, typically 15 to 20 bases long, bind to the sense mRNA or pre mRNA region coding for the protein of interest, which can inhibit translation of the bound mRNA to protein. The cDNA sequence encoding  
25 DGAT1 can thus be used to design a series of oligonucleotides which together span a large portion, or even the entire cDNA sequence. These oligonucleotides can be tested to determine which provides the greatest inhibitory effect on the expression of the protein (Stewart 1996). The most suitable mRNA target sites include 5'- and 3'- untranslated regions as well as the initiation codon. Other regions might be found to be  
30 more or less effective.

Alternatively, an antisense nucleic acid or oligonucleotide may bind to DGAT1 coding sequences.

- 35 In yet another embodiment, the invention provides a method of modulating milk production and/or milk content in a lactating bovine, including administering to the bovine an effective amount of a nucleic acid molecule having ribozyme activity and a

nucleotide sequence substantially complementary to at least a portion of mRNA encoding a bovine DGAT1 and being of sufficient length to bind selectively thereto to sufficiently reduce expression of said DGAT1.

- 5 Rather than reducing DGAT1 activity in the bovine by inhibiting gene expression at the nucleic acid level, activity of the relevant DGAT1 protein may be directly inhibited by binding to an agent, such as, for example, a suitable small molecule or a monoclonal antibody.
- 10 Thus, the invention also includes a method of inhibiting the activity of bovine DGAT1 in a lactating bovine so as to modulate milk production and/or milk solids content, comprising administering an effective amount of an antibody to the relevant DGAT1.

The invention still further includes a method of modulating milk production and/or  
15 milk solids content by raising an autoantibody to a bovine DGAT1 in the bovine. Raising the autoantibody can include administering a protein having DGAT1 activity to the bovine.

- In still a further embodiment, nucleic acids which encode DGAT1 proteins can be used  
20 to generate transgenic animals. A transgenic animal (eg. a mouse) is an animal having cells that contain a transgene, which transgene is introduced into the animal or an ancestor of the animal at a prenatal, eg. an embryonic stage. A transgene is DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, a bovine cDNA, comprising the nucleotide sequence shown in Figure 2b,  
25 or an appropriate variant or subsequence thereof, can be used to generate transgenic animals that contain cells which express the relevant DGAT1. Likewise, variants can be used to generate transgenic animals. "Knock out" animals can also be generated.

- Methods for generating transgenic animals, particularly animals such as mice, have  
30 become conventional in the art are described, for example, in US Patent Nos. 4,736,866 and 4,870,009. In such methods, plasmids containing recombinant molecules are microinjected into mouse embryos. In particular, the plasmids can be microinjected into the male pronuclei of fertilised one-cell mouse eggs; the injected eggs transferred to pseudo-pregnant foster females; and the eggs in the foster females allowed to develop to  
35 term. (Hogan, 1986). Alternatively, an embryonal stem cell can be transfected with an expression vector comprising nucleic acid encoding a DGAT1 protein, and cells containing the nucleic acid can be used to form aggregation chimeras with embryos

from a suitable recipient mouse strain. The chimeric embryos can then be implanted into a suitable pseudopregnant female mouse of the appropriate strain and the embryo brought to term. Progeny harbouring the transfected DNA in their germ cells can be used to breed uniformly transgenic mice.

5

Such animals could be used to determine whether a sequence related to an intact DGAT1 gene retains biological activity of the encoded DGAT1. Thus, for example, mice in which the murine DGAT1 gene has been knocked out and containing the nucleic acid sequence identified in Figure 2b or fragment or variant thereof could be generated. The animals could be examined with reference to milk production and content.

10

The pattern and extent of expression of a recombinant molecule of the invention in a transgenic mouse is facilitated by fusing a reporter gene to the recombinant molecule such that both genes are co-transcribed to form a polycistronic mRNA. The reporter gene can be introduced into the recombinant molecule using conventional methods such as those described in Sambrook *et al.*, (Sambrook, 1989). Efficient expression of both cistrons of the polycistronic mRNA encoding the protein of the invention and the reporter protein can be achieved by inclusion of a known internal translational initiation sequence such as that present in poliovirus mRNA. The reported gene should be under the control of the regulatory sequence of the recombinant molecule of the invention and the pattern and extent of expression of the gene encoding a protein of the invention can accordingly be determined by assaying for the phenotype of the reporter gene. Preferably the reporter gene codes for a phenotype not displayed by the host cell and the phenotype can be assayed quantitatively. Examples of suitable reporter genes include lacZ ( $\beta$ -galactosidase), neo (neomycin phosphotransferase), CAT (chloramphenicol acetyltransferase) dhfr (dihydrofolate reductase), aphIV (hygromycin phosphotransferase), lux (luciferase), uidA ( $\beta$ -glucuronidase). Preferably, the reporter gene is lacZ which codes for  $\beta$ -galactosidase.  $\beta$ -galactosidase can be assayed using the lactose analogue X-gal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside) which is broken down by  $\beta$ -galactosidase to a product that is blue in colour.

25

30

Still further transgenic applications of the invention arise from knocking out the endogenous gene encoding DGAT1 in mammals and replacing this with a transgene, in order to obtain a desired effect. This is particularly true in cattle raised for milk production. For example, additional copies of the gene encoding DGAT1 can be inserted as a transgene, or the endogenous gene associated with a high level expression promoter in a transgene. It may also prove advantageous to substitute a defective gene

35

rather than delete the entire sequence of DNA encoding for a protein having DGAT1 activity. A method of producing a transgenic bovine or transgenic bovine embryo is described in United States Patent No. 5,633,076, issued 27 May 1997, for example.

- 5 These transgenic animals of the invention can again be used to investigate the molecular basis of DGAT1 action. For example, it is expected that mutants in which one or more of the conserved cysteine residues has been deleted would have diminished activity in relation to a DGAT1 protein in which all such residues are retained. Further, deletion of a proteolytic cleavage site would likely result in a mutant lacking biological  
10 activity of DGAT1.

- Transgenic animals of the invention can also be used to test substances for the ability to prevent, slow or enhance DGAT1 activity. A transgenic animal can be treated with the substance in parallel with an untreated control transgenic animal. Substances  
15 which could be tested in this way include proteins extracted from foods ingested by the animal. For example, proteins extracted from pastoral grasses and other fodder can be tested to determine their effect on DGAT1 activity, including to determine whether breed-specific effects can be induced.

- 20 Thus, in further aspects, the invention provides transgenic non-human animals. These include by way of example only a transgenic bovine having a genome lacking a gene encoding a protein having biological activity of DGAT1 (or indeed any DGAT1 activity at all); a transgenic mouse having a genome containing a gene encoding a bovine protein having biological activity of any DGAT1; and a transgenic bovine having a gene  
25 encoding a bovine protein having biological activity of a bovine DGAT1 and heterologous nucleotide sequence antisense to the gene. The transgenic bovine can include a gene encoding a nucleic acid sequence having ribozyme activity and in transcriptional association with the nucleotide sequence antisense to the gene.

- 30 The invention further provides a transgenic bovine having a genome which includes additional copies of a gene encoding a protein having biological activity of DGAT1 or copies of a gene encoding a protein having biological activity of DGAT1 under control of a high expression promoter.

- 35 These are but a selection of the applications of this invention. Others will be apparent to those persons skilled in this art and are in no way excluded. To the contrary, the

invention extends to cover not only the specific teaching provided but also all variations and modifications which are within the skill and contemplation of the addressee.

The invention will now be defined by specific examples which are illustrative only and are not intended to limit the invention in any way.

## EXPERIMENTAL

### 1. Location of the gene responsible for the observed QTL

#### *Construction of a BAC contig spanning the BULGE9-BULGE30 interval.*

In order to clone the gene(s) responsible for the observed QTL effect, a BAC contig spanning the corresponding marker interval was constructed. This was accomplished by screening a BAC library by filter hybridisation with the microsatellite markers available for proximal BTA14q, as well as with human cDNA clones mapping to the orthologous chromosome segment on the human RH transcript map: 8q23.3-ter (Riquet *et al.*, (1999)). The ends of the isolated BACs were sequenced, sequence tagged sites (STS) developed from the corresponding sequences, and mapped onto a bovine x hamster whole genome radiation hybrid panel. This STS content mapping approach lead to the construction of the BAC contig shown in Figure 1.

#### *DGAT1 maps to the BULGE9-BULGE30 interval and is a strong positional candidate for the QTL.*

A murine gene encoding a protein with Diacylglycerol-o-acyltransferase (DGAT1) activity was identified (Cases *et al.*, (1998)) and shown to completely inhibit lactation when knocked out in the mouse (Smith *et al.*, (2000)). This gene was reported in the human to map to HSA8qter (Cases *et al.*, (1998)), ie. in the region orthologous to that containing the bovine QTL. Screening the publicly available databases with the published murine and human DGAT1 cDNA sequences allowed identification of (i) a human BAC clone containing the human DGAT1 gene (AF205589), and (ii) three bovine Expressed Sequence Tags (AW446908; AW446985; AW652329) jointly covering approximately two thirds of the bovine gene. Aligning the human DGAT1 genomic sequences with the human and bovine cDNA sequences allowed the corresponding intron-exon boundaries to be identified. Primers were developed to PCR amplify a portion of the bovine DGAT1 gene. Screening the BACs composing the BULGE9-BULGE30 contig clearly indicated that the bovine DGAT1 gene was contained in a

subset of the BACs allowing us to accurately position the DGAT1 gene in the contig of Figure 1.

These results demonstrated that the map position of DGAT1 coincided with the most likely position of the chromosome 14 QTL as determined by linkage and linkage disequilibrium analyses. Knowing that the QTL primarily affects fat content, knowing the enzymatic activity of DGAT1 and the effect of a DGAT1 knock-out on lactation, this gene was considered to be a very strong positional candidate for the corresponding QTL.

#### 10 *Organisation of the bovine DGAT1 gene*

The organisation of the bovine DGAT1 gene was determined by sequence analysis of one of the DGAT1 containing BACs. Primers were designed based on the available bovine, murine and human cDNA sequences which were either used for direct sequencing of the BAC clone or to generate PCR products corresponding to different parts of the bovine DGAT1 gene from this BAC which were then subjected to cycle-sequencing. All available sequences were then merged using the Phred / Phrap software (Ewing *et al.*, (1998); Ewing & Green, (1998); Gordon *et al.*, (1998)) to yield the consensus sequence shown in Figures 2a and b.

20 RT-PCR, 5' and 3' RACE experiments were performed on mRNA isolated from bovine mammary gland and the obtained PCR products subjected to cycle sequencing. Comparison of the genomic and cDNA sequences showed that the bovine *DGAT1* gene spans 8.6 Kb and comprises 17 exons measuring 121.8 bp on average (range: 42 – 436 bp) and allowed intron-exon boundaries to be identified (Figures 2a, 2b and 3). The cDNA sequence is also set out in SEQ ID NO: 4. While the first two introns are respectively 3.6 and 1.9 Kb long, the remaining 14 introns are only 92.4 bp long on average (range: 70 – 215 bp). All introns conform to the GT-AG rule and are strictly conserved between human and bovine. The bovine *DGAT1* gene is transcribed in a mRNA comprising >31 bp of 5' UTR sequence (Figure 2a), 1470 bp coding for a protein of 489 amino-acids, and 275 bp of 3' UTR sequence including a canonical AATAAA polyadenylation signal. The human and bovine *DGAT1* nucleotide (coding) and protein sequences are respectively 89.5% and 92.5% identical (Figures 2a, 2b, 4a and 4b). In addition, an alternative splicing variant is predicted in the bovine for exon VIII (Figure 2b). The corresponding bovine cDNAs are predicted to encode proteins comprising respectively 489 and 467 (alternative splicing variant) amino-acid residues (Figures 4a and 4b).

The predicted "Q" and "q" QTL alleles differ by a non conservative lysine to alanine amino-acid substitution in the DGAT1 gene.

Assuming that DGAT1 is indeed the QTL, it is predicted that the identified "Q" and "q" QTL alleles will correspond to functionally distinct DGAT1 alleles, ie. will differ at one or more mutations causing these alleles to be functionally different. To test this hypothesis, the structure of the DGAT1 gene in individuals predicted to be of different QTL genotypes: "QQ", "Qq" and "qq" was examined. More specifically, the DGAT1 gene from:

- (i) two sires with " $HQ^D/h^d$ " genotype as well as two of their " $HQ^D/HQ^D$ " offspring, two of their " $h^d/h^d$ " offspring and one " $HQ^D/h^d$ " offspring, and
- (ii) one " $HQ^{NZ}/h^d$ " sire with one of its " $HQ^{NZ}/HQ^{NZ}$ " offspring

was analysed wherein  $HQ^D$  corresponds to the Dutch Q haplotype and  $HQ^{NZ}$  corresponds to the New Zealand Q haplotype, and primer pairs were designed that allowed for the amplification from genomic DNA of (i) the coding portion of exon I, (ii) exon II, and (iii) the chromosome regions spanning exons III to XVII. The corresponding PCR products from the selected individuals were cycle-sequenced and the resulting sequences examined with the *Polyphred* software.

Additional sequencing analysis, as described above, on DNA from a range of breeds revealed additional polymorphisms included in Table 1 (see Methods section for breeds). Four such polymorphisms were investigated further:

(i) **K232A:** a substitution of a *ApA* by a *GpC* dinucleotide in exon VIII (respectively positions 694 and 695 counting from the start codon in the cDNA). The substitution of these two adjacent nucleotides results in a non conservative lysine (hydrophylic basic amino acid) to alanine (hydrophobic amino acid) substitution in the DGAT1 protein. The lysine residue affected by this polymorphism is conserved in the human and murine DGAT1 sequences. Together with the resulting change in the electrical charge of the protein, this strongly suggests that this amino-acid substitution is likely to result in a functional difference between the two corresponding alleles and to be at least partly responsible for the observed QTL effect.

(ii) **Nt984+8(Base 7438 A-G):** A A to G substitution in intron 12, eight base pairs downstream of exon XII. Following standard nomenclature, this polymorphism will be



referred to as *Nt984+8(A-G)*. This polymorphism cannot be predicted as such to modify the functionality of the corresponding alleles although an effect on the splicing mechanism cannot be excluded given its proximity to the intron-exon boundary.

- 5 **(iii) Nt984+26(Base 7456 C-T):** A C to T substitution in intron 12, 26 base pairs downstream of exon XII. Following standard nomenclature, this polymorphism will be referred to as *Nt984+26(Base 7456 C-T)*. Again, this polymorphism cannot be predicted as such to modify the functionality of the corresponding alleles although an effect on the splicing mechanism cannot be excluded given its proximity to the intron-exon  
10 boundary.

- (iv) Nt1470+85(Base 8402 C-T):** A C to T substitution in the 3' UTR. Following standard nomenclature, this polymorphism will be referred to as *Nt1470+85(Base 8402 C-T)*. Again, this polymorphism cannot be predicted as such to modify the functionality  
15 of the corresponding alleles although an effect on polyadenylation or mRNA stability cannot be excluded.

### Conclusion

- 20 These four polymorphisms were shown to assort into three distinct SNP haplotypes referred to as *sh<sup>Q-D</sup>*, *sh<sup>Q-NZ</sup>* and *sh<sup>a</sup>* because in the sequenced samples they coincided respectively with microsatellite haplotypes  $\mu H^{Q-D}$ ,  $\mu H^{Q-NZ}$  and  $\mu h^a$ . The base pair compositions of these three SNP haplotypes are shown in Figure 3.
- 25 Because the *sh<sup>Q-NZ</sup>* and *sh<sup>a</sup>* marker haplotypes share the G residue at the *DGAT1* *Nt984+8(Base 7438 A-G)* site, the causality of this polymorphism in the determinism of the QTL could be excluded. For the three remaining polymorphic sites, however, the *DGAT1* haplotypes associated with marker haplotypes *sh<sup>Q-D</sup>* and *sh<sup>Q-NZ</sup>* proved identical to each other while different from the *sh<sup>a</sup>* *DGAT1* haplotype. Either of these three  
30 polymorphisms could therefore be responsible for the observed QTL effect. The *Nt984+26(Base 7456 C-T)* and *Nt1470+85(Base 8402 C-T)* polymorphisms are *a priori* more likely to be neutral with respect to *DGAT1* activity because of their respective location in an intron and the 3' UTR and likewise the other non coding or neutral polymorphism shown in Table 1. A direct effect of the *K232A* mutation on *DGAT1*  
35 activity, however, is very plausible. Indeed, the corresponding lysine residue is conserved amongst all examined mammals (i.e. human, mouse, rat, pig, sheep, bison) demonstrating its functional importance (Figure 5). The evolutionary conservation of

this lysine residue also demonstrates that the *K* residue characterizing the *sH<sup>Q-D</sup>* and *sH<sup>Q-NZ</sup>* marker haplotypes is more than likely the ancestral state and that it is the *A* residue characterizing the *sh<sup>a</sup>* haplotypes that corresponds to a more recently evolved state.

5

## 2. Genotype Testing and Analysis I

This summarises the genotype testing and subsequent analysis of Holstein-Friesian animals sourced from New Zealand and Holland which were tested for the presence of the K232A polymorphism. Reference to allele "Q" corresponds to the *K* residue and allele "q" to the *A* residue (as shown in Figure 3 and Table 1).

An oligonucleotide ligation assay (OLA) was developed as described in the method section below that allows for efficient genotyping of the four *DGAT1* polymorphisms simultaneously. This OLA-test was used to genotype a previously described (Farnir et al., 2000) "grand-daughter design" (i.e. series of paternal half-brother pedigrees) comprising 1,818 Dutch Holstein-Friesian sires as well as a "daughter design" (i.e. series of paternal half-sister pedigrees) comprising 529 New Zealand Holstein-Friesian cows selected according to phenotype as described below. The marker linkage phase for each individual was determined as described below.

Fig. 6 summarizes the frequency distribution of *DGAT1* haplotypes encountered in the Dutch and New Zealand populations respectively. Four distinct *SNP* haplotypes were identified. Three of these correspond to the *sH<sup>Q-D</sup>*, *sH<sup>Q-NZ</sup>* and *sh<sup>a</sup>* that were previously identified by sequencing, and jointly account for 99% and 98% of the chromosomes in the Dutch and New-Zealand populations respectively. A fourth minor haplotype was found accounting for the remaining 1% and 2% of the chromosomes. As this haplotype codes for a *K* residue at position 232 it was assumed to correspond to a fat increasing "Q" allele and was therefore referred to as *sH<sup>Q-III</sup>* (Fig. 3). The observation that the *K* residue is found on three distinct *DGAT1* haplotypes while the *A* residue is found on a unique *DGAT1* haplotype is in agreement with *K* being the more ancient state.

The *sH<sup>Q-D</sup>* and *sH<sup>Q-NZ</sup>* *SNP* haplotypes (coding for a *K* residue at position 232) appear to be in strong linkage disequilibrium (LD) with the flanking microsatellite markers *BULGE09* and *BULGE11*, as they are in essence associated with unique microsatellite haplotypes corresponding respectively to the previously defined  $\mu H^{Q-D}$  and  $\mu H^{Q-NZ}$  haplotypes (Fig. 6C&D). In sharp contrast, the *sh<sup>a</sup>* haplotype (coding for an *A* residue at

35

position 232) is nearly evenly distributed across more than ten distinct microsatellite haplotypes (Fig. 6B).

These observations are in excellent agreement with the results of the combined linkage and LD analysis (Fernier et. al., 2000). These studies indeed predicted (i) that in the Dutch population the vast majority (estimates ranging from 81% to 92%) of "Q" allele (= K) would reside on the  $\mu H^{Q-D}$  microsatellite haplotype, (ii) that in the New Zealand population a large fraction (estimates ranging from 36% to 51%) of "Q" alleles would reside on haplotype  $\mu H^{Q-NZ}$  (we now see that the remainder correspond mainly to the  $\mu H^{Q-D}$  microsatellite haplotype) and (iii) that in both populations the "q" alleles (= A) would correspond to multiple marker haplotypes, corresponding to  $h^q$ .

Figure 7 illustrates the gain in LD signal that could be obtained in the Dutch Holstein-Friesian grand-daughter design when adding the *DGAT1* polymorphisms to the previously available markers for proximal BTA14q and performing a joint linkage and LD multipoint analysis (Fernier et. al., 2000) using the sires "daughter yield deviations" (DYD (Van Raden and Wiggans, 1991) corresponding to half breeding values) for milk fat percentage as phenotype. It can be seen that the lod score attributable to LD essentially doubles (from 3.7 to 7.8), and maximizes exactly at the position of the *DGAT1* gene. This result strongly supports the causal involvement of the *DGAT1* gene in the QTL effect. The corresponding ML estimates of the "Q" to "q" allele substitution effect ( $\alpha/2$ ) (as defined in Falconer and Mackay, 1996), residual standard deviation ( $\sigma$ ), population frequency of the "Q" allele ( $f_Q$ ), number of generations to coalescence ( $g$ ) and heterogeneity parameter ( $\rho$ ) were respectively 0.11% ( $\alpha/2$ ), 0.06% ( $\sigma$ ), 0.20 ( $f_Q$ ), 5 ( $g$ ) and 0.84 ( $\rho$ ).

Using the same Dutch Holstein-Friesian population, the additive effect of the *DGAT1* K232A polymorphism on milk yield and composition was examined. The sons DYDs for milk yield (kgs), protein yield (kgs), fat yield (kgs), protein percentage and fat percentage, were analysed using a mixed model including (i) a regression on the number of K alleles in the genotype (0, 1 or 2), and (ii) a random polygenic component estimated using an individual animal model and accounting for all known pedigree relationships. Table 2 below, reports the obtained results. It can be seen that the K232A mutation has an extremely significant effect on the five analysed dairy traits. The proportion of the trait variance explained by this polymorphism in this population ranges from 8% (protein yield) to 51% (fat percentage), corresponding to between 10% (protein yield) and 64% (fat percentage) of the genetic variance (= QTL + polygenic).

Note that the proportion of the variance explained by the full model ( $1-r^2_{\text{error}}$ ) is of the order of 70% for the yield traits and 80% for the percentage traits, which is in agreement with the known reliabilities of the corresponding DYDs (Van Raden and Wiggans, 1991). An interesting feature of this QTL effect is that the “q” to “Q” substitution increases fat yield, while decreasing milk and protein yield, despite the strong overall positive correlation characterizing the three yield traits.

Table 2:

**Effect of the *DGAT1* K232A mutation on sire's daughter yield deviations (DYDs) for milk yield and composition.**

Trait	$\alpha/2$	$r^2_{\text{QTL}}$	p-value $_{\text{QTL}}$	$r^2_{\text{polygenic}}$	$r^2_{\text{error}}$
Milk yield (Kgs)	-158 Kgs	0.18	5.00E-35	0.49	0.32
Fat yield (Kgs)	5.23 Kgs	0.15	1.57E-29	0.55	0.30
Protein yield (Kgs)	-2.82 Kgs	0.08	1.70E-15	0.65	0.26
Fat %	0.17 %	0.51	4.33E-122	0.29	0.19
Protein %	0.04 %	0.14	5.05E-28	0.66	0.20

(i)  $\alpha/2$ : QTL allele substitution effect on DYD (half breeding value), corresponding in the mixed model to the regression coefficient on the number of K alleles in the *DGAT1* K232A genotype, and to  $\alpha/2$ , where  $\alpha$  is defined according to ref. Falconer and Mackay, 1996. (ii)  $r^2_{\text{QTL}}$ : proportion of the trait variance explained by the *DGAT1* K232A polymorphism. (iii) p-value  $_{\text{QTL}}$ : statistical significance of the *DGAT1* K232A effect. (iv)  $r^2_{\text{polygenic}}$ : proportion of the trait variance explained by the random, polygenic effect in the mixed model. (v)  $r^2_{\text{error}}$ : proportion of the trait variance unexplained by the model.

The two previous analyses examined the effect of the *DGAT1* polymorphism on estimated breeding values. By definition, this phenotype will only account for the additive component of the *DGAT1* effect, and justifies the use of a regression on the number of K alleles in the mixed model. To evaluate the dominance relationship between the *DGAT1* alleles, the effect of the K232A genotype on the lactation values (first yield deviations (Van Raden and Wiggans, 1991)) of the cows composing the New Zealand daughter design were analysed. This was achieved by using a mixed model including (i) a fixed effect corresponding to the K232A genotype, and (ii) a random polygenic component accounting for all known pedigree relationships (“animal model”). Very significant effects of K232A genotype on all examined yield and composition traits were found in this population as well (Table 3, below), accounting for between 1% (protein yield) and 31% (fat percentage) of the trait variance. The observed dominance deviations,  $d$ , corresponding to the difference between the genotypic value of the KA

genotype and the midpoint between the AA and KK genotypic values (Falconer and Mackey, 1996) are shown in Table 3 below. Genotypic values of the heterozygous genotype are systematically in between alternate homozygotes. None of the  $d$ -values proved to be significantly different from zero, indicating an absence of dominance.

5 Average  $K$  to  $A$  QTL allele substitution effects,  $\alpha$  (Falconer and Mackey, 1996), were computed from the estimates of  $a$ - and  $d$ -values, as well as the population frequencies of the  $K$  and  $A$  alleles (Table 3). The predicted substitution effects are in good agreement with those computed from the grand-daughter design: the  $K$  allele increases fat yield, fat % and protein %, while decreasing milk and protein yield.

10 values of  $\alpha$  estimated from the grand-daughter and daughter design are in perfect agreement for fat and protein %, while for the yield traits estimates are larger in the grand-daughter design when compared to the daughter design. The exact reasons for this are being explored. It could be due to the fact that the sire population in the grand-daughter design is not representative of the cow population in general, or to

15 intrinsic differences between the Dutch and New-Zealand populations and/or environment.

Table 3:

**Effect of the DGAT1 K232A mutation on cows' lactation values**

**for milk yield and composition.**

20

Trait	a	d	$\alpha$	$r^2_{QTL}$	p-val $QTL$	$r^2_{polygenic}$	$r^2_{error}$
Milk yield (Kgs)	-144 Kgs	-42 Kgs	-161 Kgs	0.03	1.05E-8	0.54	0.43
Fat yield (Kgs)	7.82 Kgs	-0.89 Kgs	7.46 Kgs	0.09	1.77E-20	0.46	0.45
Protein yield (Kgs)	-2.34 Kgs	-0.76 Kgs	-2.64 Kgs	0.01	4.35E-2	0.37	0.42
Fat %	0.41 %	0.03 %	0.42%	0.31	2.5E-108	0.49	0.20
Protein %	0.08%	0.03 %	0.08%	0.04	1.60E-20	0.72	0.24

(i)  $a$ : half the difference between the genotypic values of the KK and AA genotypes (Falconer and Mackey, 1996). (ii)  $d$ : dominance deviation (Falconer and Mackey, 1996): deviation of the KA genotypic value from the midpoint between the AA and KK genotypic values; none of these proved to be significantly different from zero. (iii)  $\alpha$ : average  $K$  to  $A$  substitution effect, computed as " $a + d(q-p)$ " (Falconer and Mackey, 1996), where  $q$  is the allelic frequency of  $K$  ( $=0.7$ ) and  $p$  of  $A$  ( $=0.3$ ) (iv)  $r^2_{QTL}$ : proportion of the trait variance explained by the DGAT1 K232A polymorphism. (v) p-val  $QTL$ : statistical significance of the DGAT1 K232A effect. (vi)  $r^2_{polygenic}$ : proportion of the trait variance explained by the random, polygenic effect in the mixed model. (vii)  $r^2_{error}$ : proportion of the trait variance unexplained by the model.

30

*Pedigree material and phenotypes.* The pedigree material used for the association studies comprised a "grand-daughter design" (Weller et. al., 1990) counting 1,818 Holstein-Friesian bulls sampled in the Netherlands, as well as a "daughter-design" (Weller et. al., 1990) counting 529 Holstein-Friesian cows sampled in New Zealand. The phenotypes of the sires were "daughter yield deviations" (DYD: unregressed weighted averages of the daughter's lactation performances adjusted for systematic environmental effects and breeding values of the daughter's dams and expressed as deviations from the population mean (Van Raden and Wiggans, 1991)) obtained directly from CR-Delta (Arnhem - The Netherlands). The phenotypes of the cows were "lactation values" (first lactation yield deviations (YD), i.e. weighted average lactation performances expressed as deviations from the population mean, adjusted for management group, permanent environmental effects and herd-sire interaction effects (Van Raden and Wiggans, 1991)) obtained directly from Livestock Improvement Corporation (Hamilton - New Zealand).

*Combined linkage and linkage disequilibrium analysis and association studies.* The maximum likelihood procedure for combined linkage and linkage disequilibrium analysis is described in detail in Farnir, 2000. The association study in the grand-daughter design was performed using the following model:

$$y_i = \mu + \beta x_i + a_i + e_i$$

where  $y_i$  is the DYD of son  $i$ ,  $\mu$  is the overall population mean,  $\beta$  is a fixed regression coefficient estimating the A to K allele substitution effect,  $x_i$  is an indicator variable corresponding to the number of K alleles in the K232A genotype,  $a_i$  is a random polygenic component accounting for all known pedigree relationships ("animal model" Lynch and Walsh, 1997) and  $e_i$  is a random residual. The association study in the daughter design was performed using the model:

$$y_i = \mu + g_i + a_i + e_i$$

where  $y_i$  is the lactation value of cow  $i$ ,  $g_i$  is a fixed effect corresponding to the DGAT1 genotype (KK, KA, or AA),  $a_i$  is a random polygenic component accounting for all known pedigree relationships ("animal model" Lynch and Walsh, 1997) and  $e_i$  is a random residual. In both instances, maximum likelihood solutions for  $\beta$ ,  $g_i$ ,  $a_i$ ,  $e_i$ ,  $\sigma^2_a$ ,  $\sigma^2_e$  were obtained using the MTDFREML program (Boldman et al, 1997).

### 3. Genotype Testing and Analysis II

This summarises the genotype testing and subsequent analysis of Holstein-Friesian,  
 5 Jersey and Ayrshire animals in a separate population from those presented in genotype  
 testing and analysis I, above.

#### *Progeny tested sires*

Each year Livestock Improvement Corporation (New Zealand) progeny test some 200-  
 10 300 bulls per year. This entails the bulls being genetically evaluated on the basis of 50-  
 85 daughters per sire. The sires are evaluated for milk fat, milk protein, milk volume  
 and 20 non-production traits. Semen has been retained from all progeny tested sires  
 since the early 1970s. DNA was extracted from the semen and genotyped for the K232A  
 DGAT1 polymorphism using the 7900 Taqman system (see Methods section below).

15 Statistical analysis was undertaken on this dataset using Restricted Maximum  
 Likelihood (REML) and the average information algorithm (Johnson and Thompson,  
 1995). The linear model included the fixed effects of DGAT1 (3 classes; 0, 1 and 2  
 copies of the Q allele i.e. the K residue) and a covariate corresponding to the proportion  
 20 of overseas genetics. The random effect was animal with a relationship matrix based on  
 all known relationships. Daughter yield deviations (DYDs), weighted averages of a sire's  
 daughter's lactation performances expressed as deviations from the population mean  
 (van Raden and Wiggans 1991) were used as the phenotypic measurement. The  
 phenotypes were weighted by a weighting factor based on the variance of the DYD for a  
 25 son being:

$$\text{Var DYD} = \left[ \frac{1 + (n-1)\frac{1}{4}h^2}{n} \right] \sigma_p^2$$

where Var DYD is the variance of son's DYD; n is the number of daughters contributing  
 30 to the DYD; h<sup>2</sup> is the heritability, which was taken as 0.35 for yield traits.

The dataset was analysed separately for the 3 major breeds; Holstein-Friesian, Jersey  
 and Ayrshire.

Seventeen hundred and thirteen Holstein-Friesian sires were included in the analysis. The effect of the DGAT1 polymorphism was extremely significant for the three milk production traits (Table 4). With each additional Q allele the level of milk fat production increases by approximately 6 kg per lactation, milk protein production decreases by approximately 2.5 kg per lactation and milk volume decreases by approximately 125 litres per lactation.

Table 4:

**Effect of the DGAT1 polymorphism on milk production in the Holstein-Friesian bull population (kilograms per lactation).**

	Fat	Protein	Milk
qq	0	0	0
Qq	6.86	-2.13	-128
QQ	11.83	-4.80	-266
st. error	0.87	0.68	24

The effects for the Jersey and Ayrshire breeds were less significant than those of the Holstein-Friesian breed but were consistent in direction of effects.

#### *Daughters for milk components*

Data collection was integrated with LIC's herd testing service using a sample of 102 herds involved in LIC's Sire Proving Scheme (SPS) in 1995. In addition to milk volume from herd testing, the concentrations of fat, crude protein (total nitrogen), casein, whey and lactose were determined. The data was collected from over 3,000 cows born in 1996 and first calving in the 1998 spring season, these being predominantly the daughters of approximately 220 SPS bulls. The milk characteristics were measured at three herd tests on each cow, with each herd having a herd test in each of the Sept/Oct, Nov/Dec and Jan/Feb periods. The Milkoscan FT120, which employs Fourier transform infrared spectrophotometry with enhanced milk calibrations (Foss Electric Application Note Nos. 95, P/N 492280 and 102, P/N 578377), was used to determine the milk component concentrations.

Nine hundred and twelve daughters were genotyped for the DGAT1 polymorphism using the OLA system. Analysis was undertaken using SAS (Statistics, Version 5, 1985) fitting a general linear model. The model included sire and maternal grandsire as fixed effects, DGAT1 polymorphism (3 classes; 0, 1 and 2 copies of the Q allele), covariates including 16ths of Holstein-Friesian, Jersey, Ayrshire and other, proportion of overseas



genetics within the Holstein-Friesian, Jersey and Ayrshire breeds. Yield deviations that were pre-adjusted for herd, stage of lactation among other fixed effects were used (Johnson et al 2000).

- 5 The DGAT1 polymorphism is statistically significant for Lactose, casein, beta-casein and whey yield and also for casein and beta-casein percent as outlined in Table 5.

**Table 5:**  
**Effect of the DGAT1 polymorphism on milk components.**

Trait	qq	Qq	QQ	p-value
Lactose yield*	48	23	0	<0.0001
Casein yield*	11.0	5.8	0	0.01
Casein %	-0.13	-0.06	0	<0.0001
Whey yield*	6.86	2.31	0	<0.0001
$\beta$ -casein yield*	3.98	2.19	0	0.05
$\beta$ -casein %	-0.43	-0.23	0	0.0001

\* Units = g/day for lactose, casein and whey yield and g/litre for  $\beta$ -casein yield

#### *Daughters for solid fat content*

- 15 Six hundred and ninety-two daughters were phenotyped for solid fat content. Solid fat content of the milkfat is a characteristic which has a major influence on the functionality of milkfat products, and in particular has a significant effect on the hardness of butter (MacGibbon & McLennan, 1987). The solid fat content at 10°C (SFC 10) was used for comparison of the properties of the milkfat as it relates well to the
- 20 sectility hardness measurement of butter, a major functional property. Thus the performance of milkfat products may be predicted from the characteristics of the milk produced. The solid fat content (SFC) of the extracted fat was determined by pulsed nuclear magnetic resonance (NMR) and expressed as percentage solid fat (MacGibbon & McLennan, 1987). As the milkfat was melted to remove any thermal history, prior to
- 25 recrystallization under standard conditions, the SFC simply reflects the chemical composition of the milkfat.

The 692 daughters were a subset of the 912 daughters that were phenotyped and genotyped for the results presented in Table 2. The solid fat content measures were

collected over 2 lactations. Breeding values were calculated using an animal model similar to that of Johnson et al 2000.

- 5 The same statistical model was fitted for solid fat content as was for the milk component analysis. The DGAT1 polymorphism has a statistically significant effect (p-value <0.0001) on solid fat content, increasing it by approximately by 1% for each addition of the Q allele.

- 10 This effect was further confirmed in 50 daughters (predominantly Holstein-Friesian) that were farmed at one location and measured for SFC on the same day. The estimated effect for of the DGAT1 polymorphism on SFC was to increase it by approximately 2% per addition of each Q allele. This finding was significant at the five percent threshold level.

- 15 The genetic standard deviation for SFC is 2.25 (D Johnson personal communication) and thus the effect of DGAT1 is approximately 0.5 of a genetic standard deviation.

#### **4. Relative Transcript Levels of the Splice Variant**

- 20 Real time PCR experiments were conducted using reverse transcribed mRNA isolated from lactating bovine mammary gland(s) (see experimental methods). These experiments revealed that the alternatively spliced transcript as shown on Figure 2b, was approximately 100 fold less abundant than the full length transcript.

## METHODS SECTION

In order to identify other polymorphisms within the bovine DGAT1 gene, DNA was isolated from sperm, PCR amplified and then using primers designed from the sequence shown in Figures 2a and 2b and/or the cDNA sequence (SEQ ID NO: 4) direct sequenced on an ABI 3100. The breeds examined were:

Ayrshire, Angler, Belgian Blue, Blond D'Aquitaine, Brown Swiss, Charolais, Red Devon, Devon, Dexter, Friesian, Guernsey, Belted Galloway, Gelbvieh, Hereford, Jersey, Limousin, Longhorn, Maine Anjou, MRI (Meuse-rhine-ysse), Murray Grey, Piedmontese, Romangola, Sahiwal, Santa Gertrudis, Scottish Highland, Shorthorn, South Devon, Sussex, Swedish Red, Simmental, Wagyu, Welsh Black, Angus, and Zebu.

All the polymorphisms discovered are listed in Table 1, above.

The majority of the primers are also listed in Figure 2b or contained in the cDNA sequence (SEQ ID NO: 4).

### Experimental method for the OLA analysis of four SNP's in DGAT1

#### *PCR amplification of the regions containing the polymorphisms*

Protocol for the PCR amplification of exon VIII, intron XII and 3' UTR, the regions containing the four polymorphisms that were initially described in the DGAT1 gene.

Component	For 1 sample	Final concentration
HotStar Qiagen Buffer (10 x)	1.5 µl	0.7 µM
Primer 17F at 100µM	0.07 µl	0.7µM
18R at 10 µM	0.07 µl	0.5 µM
Primer 6F at 100µM	0.05µl	0.5 µM
AW 446985dn1 at 100µM	0.05µl	0.5 µM
Primer InsUp1	0.05µl	0.5 µM
14R2	0.05µl	10%
DMSO	1 µl	300 µM
dNTP 10mM	0.3 µl	0.1 U/µl
HotStarQiagen Taq (CatNr 203205:5U/µl)	0.2 µl	
H2O	1.66 µl	
DNA (5 ng/µl)	5 µl	
Total	10 µl	

Primer sequences are given in the following table as well as the genomic region targeted by them.

SNP targeted	Primer name	Primer sequence	5' base position
Exon VIII SNP (DG 1)	17F	CCTGAGCTTGCTCTCCACAGT	6579
	18R	CCAGGAGTCGCCGAGCAGGAAG	7058
Exon XII SNPs (DG 2 and DG3)	6F	CCGGCCATCCAGAACTCCATGAAG	7280
	AW446985 dn1	TAGAACTCGCGGTCTCCAAAC	7605
	InsUp1	TGGCTGTCACTCATCATCGGGCA	8222
3'UTR SNP (DG4)	14R2	TTGCACAGCACTTTATTGACACA	8566

PCR amplification was performed on MJ PTC100 or PTCT200 cyclers using the following steps:

Step	Temperature	Time	Comment
1° Activation of the enzyme	94 °C	12 minutes	One times
2° Denaturation	92 °C	1 minute	Repeat step 2 to 4, 35 times
3° Hybridisation	60°C	1 minute 30 seconds	
4° Elongation	72°C	1 minute 30 seconds	
5° Inactivation of the enzyme	99°C	45 minutes	

5

#### Oligonucleotide Ligation Assay (OLA)

The oligonucleotides used in the OLA multiplex reaction are given in the table below. The detection of each mutation relies on the use of two fluorescent-labelled oligonucleotide (SNPx\_FAM and SNPx\_HEX) and one common 3' and 5' phosphorylated, non-labelled oligonucleotide (SNPx\_2P)

10

Locus	Oligo	Sequence	5' base position	Number of spacer phosphoramidites	Size of the ligation product*
DG1	SNP1_FAM	AGC TTT GGC AGG TAA GGC	6813	0	32
	SNP1_HEX	AGC TTT GGC AGG TAA GAA	6813		
	SNP1_2P	GGC CAA CGG GGG AG	6831		
DG2	SNP2_FAM	GCT GGC GGT GAG TGA	7424	3	39
	SNP2_HEX	GCT GGC GGT GAG TGG	7424		
	SNP2_2P	CCT GCT GGG TGG GGA	7439		
DG3	SNP3_FAM	GCT GGG TGG GGA CGC	7442	0	29
	SNP3_HEX	GCT GGG TGG GGA CGT	7442		
	SNP3_P	GTG GGG GCG GGT GG	7457		
DG4	SNP4_FAM	TGC CCC AAC CTG GGT	8388	2	36
	SNP4_HEX	TGC CCC AAC CTG GGC	8388		
	SNP4_2P	GCA GCA GGA GGA GGC	8403		

\* The size of the ligation products is the sum of the number of nucleotides of the two ligated oligonucleotides plus 3 bases equivalents per spacer phosphoramidites molecule, present at the 5' end of the common oligonucleotide.

15

For each SNP a mixture of the three oligonucleotides was prepared first, following the dilution guidelines in the table below.

SNP mixture	Oligonucleotide to mix	Quantity	Final concentration
DG1 (oligo. mixture)	SNP1_FAM 10 $\mu$ M	10 $\mu$ l	1 $\mu$ M
	SNP1_HEX 10 $\mu$ M	20 $\mu$ l	2 $\mu$ M
	SNP1_2P 10 $\mu$ M	20 $\mu$ l	2 $\mu$ M
	H <sub>2</sub> O	50 $\mu$ l	
DG2 (oligo. mixture)	SNP2_FAM 10 $\mu$ M	10 $\mu$ l	1 $\mu$ M
	SNP2_HEX 10 $\mu$ M	20 $\mu$ l	2 $\mu$ M
	SNP2_2P 10 $\mu$ M	20 $\mu$ l	2 $\mu$ M
	H <sub>2</sub> O	50 $\mu$ l	
DG3 (oligo. mixture)	SNP3_FAM 10 $\mu$ M	10 $\mu$ l	1 $\mu$ M
	SNP3_HEX 10 $\mu$ M	20 $\mu$ l	2 $\mu$ M
	SNP3_2P 10 $\mu$ M	20 $\mu$ l	2 $\mu$ M
	H <sub>2</sub> O	50 $\mu$ l	
DG4 (oligo. mixture)	SNP4_FAM 10 $\mu$ M	10 $\mu$ l	1 $\mu$ M
	SNP4_HEX 10 $\mu$ M	30 $\mu$ l	3 $\mu$ M
	SNP4_2P 10 $\mu$ M	20 $\mu$ l	2 $\mu$ M
	H <sub>2</sub> O	40 $\mu$ l	

The ligation reaction for one sample was performed as follow:

Component	Quantity persample
DG1 oligonucleotide mixture* (35, 70 and 70 nM)	0.7 $\mu$ l
DG2 oligonucleotide mixture* (12.5, 25 and 25 nM)	0.25 $\mu$ l
	0.25 $\mu$ l
DG3 oligonucleotide mixture* (12.5, 25 and 25 nM)	0.25 $\mu$ l
	2 $\mu$ l
	2 $\mu$ l
DG4 oligonucleotide mixture* (12.5, 37.5 and 25 nM )	
DMSO	1 $\mu$ l
Incubation buffer of the Tsc DNA ligase (Roche, Cat Nr 1 939 807 or 1 939 815)	8.55 $\mu$ l
Tsc DNA ligase	5 $\mu$ l
H <sub>2</sub> O	
Multiplex PCR (see above)	
Total	20 $\mu$ l

5

\* The final concentration of the oligonucleotides in the ligation reaction is given between parenthesis (SNPx\_FAM, SNP<sub>x</sub>\_HEX and SNP<sub>x</sub>\_2P respectively)

10 The sample was submitted to the following temperature cycling program in a MJ PTC100 or PTC 200 PCR machine.

Step	Temperature	Time	Comment
1° Initial denaturation step	98 °C	2 minutes	One times
2° Denaturation	94 °C	30 seconds	Repeat step 2 to 3, 30 times
3° Hybridisation and ligation	45°C	3 minutes	
5° Inactivation of the enzyme	99°C	45 minutes	

15 Following the LCR, 20  $\mu$ l of H<sub>2</sub>O was added to the ligation reaction. To 0.5  $\mu$ l of the diluted ligation reaction, either 2  $\mu$ l of loading buffer was added, or 2  $\mu$ l loading buffer containing TAMRA350 internal line size standard.

The loading buffer was composed as follows: 1 part of blue dextran (50mg/ml)/ EDTA (25mM) and 6 parts of formamide

The TAMRA350 containing loading buffer was composed as follows: 3 parts TAMRA350 (Applied Biosystems 401736 ; 8nM), 10 parts of Blue dextran (50 mg/ml)/EDTA (25 mM) and 60 parts of formamide.

- 5 TAMRA containing samples was placed alternately with TAMRA free samples when loaded onto the sequencing gel, in order to ease the identification of the lanes on the gel image.

- 10 The samples may require further dilution in order to avoid a too intense fluorescent signal on the sequencer. It is also very likely that from one primer batch to another, oligonucleotides concentrations will need adjustment.

- 15 The samples were denatured for 5 minutes at 95°C before loading. The samples were then loaded onto a 6% denaturing acrylamide gel on sequencer ABI 373 or a 4% gel on sequencer ABI 377.

In addition to the OLA assays referred to above, genotyping of the DGAT1 polymorphism was carried out by utilizing two different techniques for detection of PCR products.

*Gel-based Genotyping Assay*

- 20 Primer sequences 5' to 3', genomic sequence position in brackets:

DGAT1 21: GTAGCTTTGGCAGGTAAGAA (6811)

DGAT1 22: GGGGCGAAGAGGAAGTAGTA (6984)

DGAT1 23: TGGCCCTGATGGTCTACACC (6613)

DGAT1 24B: GGGCAGCTCCCCCGTTGGCCGC (6850)

25

- The final reaction conditions were 1X Gold PCR buffer, 2.5mM MgCl<sub>2</sub> (Applied Biosystems), 200μM each dNTP (Roche), 600nM DGAT1 21 and DGAT1 22, 400nM DGAT1 23 and DGAT1 24B (Invitrogen), 10% dimethylsulphoxide (Sigma), 3μl DNA template and 2.5 units AmpliTaq Gold DNA polymerase (Applied Biosystems) in a total  
30 volume of 50μl.

- Cycling conditions were a 94°C initial denaturation for 5 minutes, then 35 cycles of denaturation at 94°C for 30 seconds, annealing at 56°C for 30 seconds, extension at 72°C for 20 seconds followed by one extension cycle of 72°C for 2 minutes.

35

Primer positions around polymorphism (in bold) on genomic sequence from 6587 to 6986.

```

                    DGAT123   TGGC  CCTGATGGTC TACACC
                                ──────────▶
5  TGCCTCTCCC ACAGTGGGCT CCGTGCTGGC CCTGATGGTC TACACCATCC

TCTTCCTCAA GCTGTTCTCC TACCGGGACG TCAACCTCTG GTGCCGAGAG

10  CGCAGGGCTG GGGCCAAGGC CAAGGCTGGT GAGGGCTGCC TCGGGCTGGG

GCCACTGGGC TGCCACTTGC CTCGGGACCG GCAGGGGCTC GGCTCACCCC
15
                    DGAT1 21 GTAGCT TTGGCAGGTA AGAA
                                ──────────▶
CGACCCGCCC CCTGCCGCTT GCTCGTAGCT TTGGCAGGTA AGAAGGCCAA
                                ◀────────
                                CGCCGGTT

20  CGGGGGAGCT GCCCAGCGCA CCGTGAGCTA CCCCAGACAAC CTGACCTACC
GCCCCCTCGA CGGG DGAT1 24B

GCGGTGAGGA TCCTGCCGGG GGCTGGGGGG ACTGCCCCGC GGCCTGGCCT

25  GCTAGCCCCG CCCTCCCTTC CAGATCTCTA CTACTTCCTC TTCGCCCCCA
                                ◀────────
                                AT GATGAAGGAG AAGCGGGG DGAT1 22

```

30 The Q allele has polymorphic sequence AA and is detected by the DGAT1 21 + 22 primers, producing a band of 174bp. The q allele has polymorphic sequence GC and is detected by the DGAT123 + 24 primers, producing a band of 238bp.

35 The primers DGAT123 and DGAT122 also successfully PCR the DGAT1 gene producing a product of 372bp in all reactions. Therefore, a QQ homozygote would have bands at 372bp and 174bp, a qq homozygote would have bands at 372bp and 238bp and a Qq heterozygote would have all 3 bands at 372bp, 238bp and 174bp.

18µl of PCR product was separated on a 1.2% agarose TAE gel, stained with ethidium bromide and scored independently by two investigators on the basis of the number and size of bands present.

*TaqMan Allelic Discrimination Genotyping Assay*

- 5 Primer sequences 5' to 3', genomic sequence position in brackets:

DGAT1forAD: TTCTCCTACCGGGACGTCAA (6651)

ReverseNZ: CCGCGGTAGGTCAGGTTGTC (6890)

Probe sequences 5' to 3', genomic sequence position in brackets:

- 10 ForAA (FAM): CGTTGGCCTTCTTA (6838)

DGAT1ADGC (VIC): TTGGCCGCCTTACC (6836)

Both probes use MGB (minor groove binder) as a non-fluorescent quencher.

- The final reaction conditions are 1x Universal PCR Mastermix (Applied Biosystems),  
15 500nM each primer (Invitrogen), 70nM ForAA (FAM) probe, 300nM DGAT1ADGC (VIC) probe (Applied Biosystems) and 2µl of a 1/20 dilution of DNA template in a total volume of 10µl.

- Cycling conditions were 50°C for 2 minutes, 95°C initial denaturation for 10 minutes, then 37 cycles of denaturation at 94°C for 15 seconds, annealing and extension 60°C  
20 for 1 minute.

Primer positions around polymorphism (in bold) on genomic sequence from 6587 to 6986.

- 25 TGCCTCTCCC ACAGTGGGCT CCGTGCTGGC CCTGATGGTC TACACCATCC

DGAT1forAD TTCTCC TACCGGGACG TCAA →  
TCTTCCTCAA GCTGTTCTCC TACCGGGACG TCAACCTCTG GTGCCGAGAG

- 30 CGCAGGGCTG GGGCCAAGGC CAAGGCTGGT GAGGGCTGCC TCGGGCTGGG

- GCCACTGGGC TGCCACTTGC CTCGGGACCG GCAGGGGCTC GGCTCACCCC  
35



40

**ForAA(FAM) A T TCTTCCGGTTGC**

CGACCCGCCC CCTGCCGCTT GCTCGTAGCT TTGGCAGGTA AGAAGGCCAA

**DGAT1ADGC (VIC) CCAT TCCGCCGGTT**

5 CGGGGGAGCT GCCCAGCGCA CCGTGAGCTA CCCCAGACAAC CTGACCTACC  
 ← CTGTTG GACTGGATGG

GCGGTGAGGA TCCTGCCGGG GGCTGGGGGG ACTGCCCCGGC GGCCTGGCCT  
 CGCC ReverseNZ

10

GCTAGCCCCG CCCTCCCTTC CAGATCTCTA CTACTTCCTC TTCGCCCCCA

A 240bp product is produced in this reaction. When the Q allele (AA) is present the FAM-labelled probe binds and fluoresces at 518nm. When the q allele (GC) is present  
 15 the VIC-labelled probe binds and fluoresces at 554nm. After cycling is complete, the plate is scanned on the ABI7900 Sequence Detection System, the fluorescence from each well detected, and a scattergraph is drawn. The scattergraph separates out into 3 clumps with Q homozygotes in the upper left hand corner, q homozygotes in the lower right hand corner and Qq heterozygotes in between. Each clump is circled and the  
 20 software automatically determines the genotype for each sample. On each plate there are controls with 8 wells each of known Q homozygotes, q homozygotes, Qq heterozygotes and no template controls.

#### *Splice Variant Gene Expression*

25 To determine the relative gene expression of the splice variants created by insertion/deletion of 66bp around the polymorphic site by alternate exon usage, RNA was extracted from mammary tissue and reverse transcribed using oligodT primer using a first strand cDNA synthesis kit (Invitrogen). Real time PCR to determine relative quantities of each variant was then carried out.

30

Primer sequences 5' to 3', genomic sequence position in brackets:

DGAT1forRT66: TCTCCTACCGGGACGTCAAC (6652)

DGAT1revRT66: GAGATCGCGGTAGGTCAGGTT (6964)

DGAT1forRTless66: GCTGCTTTGGCAGATCTCTACTACTT (6711)

35 DGAT1revRTless66: AAGCGCTTTCGGATGCG (7038)

Probe sequences 5' to 3', genomic sequence position in brackets:

DGAT1with66 (FAM): CCGTGAGCTACCC (6857)

DGAT1less66 (VIC): CTTGCCCCCACCCT (6976)

- 5 Both probes use MGB (minor groove binder) as a non-fluorescent quencher.

Final reaction conditions were 1X Universal PCR Mastermix (Applied Biosystems), 60nM each primer (Invitrogen), 60nM each probe (Applied Biosystems) and 1µl of template cDNA in a total volume of 10µl.

- 10 Cycling conditions were 50°C for 2 minutes, 95°C initial denaturation for 10 minutes, then 37 cycles of denaturation at 94°C for 15 seconds, annealing and extension 60°C for 1 minute.

- 15 Primer positions around 66bp insertion (in italics) on cDNA sequence. The start of the cDNA sequence is equivalent to position 6479 on the genomic sequence, with the last base of the cDNA equivalent to position 7428 of the genomic sequence.

CCGTGGCCTT TCTCCTCGAG TCTATCACTC CAGTGGGCTC CGTGCTGGCC

- 20 DGAT1forRT66 TCTCCT ACCGGGACGT  
CTGATGGTCT ACACCATCCT CTTCTCAAG CTGTTCTCCT ACCGGGACGT

CAAC → DGAT1forRTless66 GCTGCTT  
CAACCTCTGG TGCCGAGAGC GCAGGGCTGG GGCCAAGGCC AAGGCTGCTT

- 25 TGGCAG DGAT1with66(FAM) C CGTGAGCTAC  
TGGCAGGTAA *GAAGGCCAAC* GGGGGAGCTG CCCAGCGCAC CGTGAGCTAC

- CC ATCTCTAC TACTT →  
30 CCCGACAACC TGACCTACCG CGATCTCTAC TACTTCCTCT TCGCCCCAC  
← TTGG ACTGGATGGC GCTAGAGDGAT1revRT66 CT TCGCCCCAC

- CCTGTGCTAC GAGCTCAACT TCCCCGCTC CCCCCGCATC CGAAAGCGCT  
CCT DGAT1less66 (VIC) ← GCGTAG GCTTTCGCGA

- 35 TCCTGCTGCG GCGACTCCTG GAGATGCTGT TCCTACCCA GCTCCAGGTG  
A DGAT1revRTless66

GGGCTGATCC AGCAGTGGAT GGTCCCGGCC ATCCAGAACT CCATGAAGCC

5 CTCAAGGAC ATGGACTACT CCCGCATCGT GGAGCGCCTC CTGAAGCTGG

This reaction detects the presence of the insertion splice variant by creating a 145bp product which binds the FAM probe only. The deletion splice variant is detected by a 92bp product that binds the VIC probe only.

10

The cDNA for each alternate splice variant was cloned into pGemT (Promega). A dilution series of the same, known amount, of each variant plasmid DNA was used to create a standard curve that established the linearity of the PCR reaction over a range of DNA concentrations. The threshold cycle number of the sample variants was  
15 converted back to a DNA amount by linear regression and the amounts of each variant present compared.

The presence of an alternate splice variant raises the possibility of an alternate function that is at this stage unknown.

20

It will be appreciated that it is not intended to limit the invention to the above examples only, many variations, which may readily occur to a person skilled in the art, being possible without departing from the scope thereof as defined in the accompanying claims.

25

#### **INDUSTRIAL APPLICATION**

The present invention is directed to a method of genotyping bovine for improved milk production traits. In particular, such traits include increased milk volume and milk  
30 protein content and decreased milkfat content and solid fat content. It is anticipated that herds of bovine selected for such a trait will produce milk which will be more easily processed and such milk and products made therefrom may provide health benefits to consumers, as well as producing an increased milk yield.

## REFERENCES

- CASES, S.; SMITH, S.J.; ZHENG, Y.-W.; MYERS, H.M.; LEAR, S.R.; SANDE, E.; NOVAK, S.; COLLINS, C.; WELCH, C.B.; LUSIS, A.J.; ERICKSON, S.K.; FARESE, R.V. JR (1998). Identification of a gene encoding an acyl CoA:diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis. *Proceedings of the National Academy of Sciences, USA* 95(22):13018-23.
- 10 COPPIETERS, W.; RIQUET, J.; ARRANZ, J.-J.; BERZI, P.; CAMBISANO, N.; GRISART, B.; KARIM, L.; MARCQ, F.; SIMON, P.; VANMANSHOVEN, P.; WAGENAAR, D.; GEORGES, M. (1998) A QTL with major effect on milk yield and composition maps to bovine chromosome 14. *Mammalian Genome* 9: 540-544.
- 15 DELOUKAS et al (1998) A physical map of the 30,000 human genes. *Science* 282, 744-746
- EWING, B.; HILLIER, L.; WENDL, M.C.; GREEN, P. (1998). Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Research* 8(3):175-185.
- 20 EWING, B.; GREEN, P. (1998). Base-calling of automated sequencer traces using Phred. II. Error probabilities. *Genome Research* 8(3):186-194.
- GORDON, D.; ABAJIAN, C.; GREEN, P. (1998). Consed: a graphical tool for sequence finishing. *Genome Research* 8(3): 195-202.
- 25 RIQUET, J.; COPPIETERS, W.; CAMBISANO, N.; ARRANZ, J.-J.; BERZI, P.; DAVIS, S.; GRISART, B.; FARNIR, F.; KARIM, L.; MNI, M.; SIMON, P.; TAYLOR, J.; VANMANSHOVEN, P.; WAGENAAR, D.; WOMACK, J.E.; GEORGES, M. (1999). Identity-by-descent fine-mapping of QTL in outbred populations: application to milk production in dairy cattle. *Proceedings of the National Academy of Sciences, USA* 96: 9252-9257.
- 30 FARNIR, F.; GRISART, B.; COPPIETERS, W.; RIQUET, J.; BERZI, P.; CAMBISANO, N.; KARIM, L.; MNI, M.; SIMON, P.; WAGENAAR, D.; GEORGES, M. (2000). Simultaneous mining of linkage and linkage disequilibrium to fine-map QTL in outbred half-sib pedigrees: revisiting the location of a QTL with major effect on milk production on bovine chromosome 14. *Ph.D Thesis, University of Liege* 2000.
- 35

- SMITH, S.J.; CASES, S.; JENSEN, D.R.; CHEN, H.C.; SANDE, E.; TOW, B.; SANAN, D.A.; RABER, J; ECKEL, R.H.; FARESE, R.V. JR (2000). Obesity resistance and multiple mechanisms of triglyceride synthesis in mice lacking DGAT. *Nature Genetics* 5 25(1):87-90.
- BARENDSE, W.; ARMITAGE, S.M.; KOSSAREK, L.M.; SHALOM, A.; KIRKPATRICK, B.W.; RYAN, A.M.; CLAYTON, D.; LI, L.; NEIBERGS, H.L.; ZHANG, N.; GROSSE, W.M.; WEISS, J.; CREIGHTON, P.; MCCARTHY, F.; RON, M.; TEALE, A.J.; FRIES, R.; 10 McGRAW, R.A.; MOORE, S.S.; GEORGES, M.; SOLLER, M.; WOMACK, J.E.; HETZEL, D.J.S. (1994). A genetic linkage map of the bovine genome. *Nature Genet.* 6: 227-235.
- BISHOP, M.D.; KAPPES, S.M.; KEELE, J.W.; STONE, R.T.; SUNDEN, S.L.F.; HAWKINS, G.A.; SOLINAS TOLDO, S.; FRIES, R.; GROSZ, M.D.; YOO, J.; BEATTIE, C.W. (1994). A 15 genetic linkage map for cattle. *Genetics* 136: 619-639.
- COLLINS, F.S. (1995). Positional cloning moves from perditional to traditional. *Nature Genet.* 9: 347-350.
- 20 GEORGES, M.; ANDERSSON, L. (1996). Livestock genomics comes of age. *Genome Research.* 6: 907-921.
- GEORGES, M; NIELSEN, D.; MACKINNON, M.; MISHRA, A.; OKIMOTO, R.; PASQUINO, A.T.; SARGEANT, L.S.; SORENSEN, A.; STEELE, M.R.; ZHAO, X.; WOMACK, J.E.; 25 HOESCHELE, I. (1995). Mapping quantitative trait loci controlling milk production by exploiting progeny testing. *Genetics* 139: 907-920.
- HOUBENWCYL (1987). *Methods of Organic Chemistry*, ed. E. Wansch. Vol. 15 I and II. Thieme, Stuttgart.
- 30 HUSE *et al.* (1989). *Science* 246: 1275-1281.
- KAPPES, S.M.; KEELE, J.W.; STONE, R.T.; McGRAW, R.A.; SONSTEGARD, T.S.; SMITH, T.P.L.; LOPEZ-CORRALES, N.L.; BEATTIE, C.W. (1997) A Second-Generation 35 Linkage Map of the Bovine Genome. *Genome Research* 7: 235-249.
- KOHLER and MILSTEIN (1975). *Nature.* 256: 495-497.

- KOZBOR *et al.* (1983). *Immunol. Today* **4**: 72.
- MERRIFIELD (1964). *J. Am. Chem. Assoc.* **85**: 2149-2154.
- 5 McCAFFERTY *et al.* (1990) *Nature* **348**: 552-554.
- SAMBROOK, J.; FRITSCH, E.F.; MANIATIS, T. (1989). *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbour Lab Press, Cold Spring Harbour, New York.
- 10 STEWART, A.J.; CANITROT, Y.; BARACCHINI, E.; DEAN, N.M.; DEELEY, R.G. and COLE, S.P.C. (1996). Reduction of expression of the multidrug resistance protein (MRP) in human tumour cells by antisense phosphorothioate oligonucleotides. *Biochem. Pharmacol.* **51**: 461-469.
- 15 VAN RADEN, P. M., AND G. R. WIGGANS, 1991 Derivation, calculation, and use of National Animal Model Information. *J. Dairy Sci.* **74**: 2737-2746.
- JOHNSON, D.L.; THOMPSON, R. 1995. Restricted maximum likelihood estimation of variance components for univariate animal models using sparse matrix techniques and average information. *J. Dairy Sci.* **78**: 449-456.
- 20 JOHNSON D.L., S.F. PETCH, A.M. WINKELMAN AND M. BRYANT 2000 Genetics of milk characteristics in New Zealand dairy cattle Proceedings of New Zealand Society of Animal Production **60**:318-319.
- 25 MACGIBBON A.K.H. & MCLENNAN W. D. 1987. Hardness of New Zealand patted butter: seasonal and regional variations. *New Zealand Journal of Dairy Science and Technology*, **22**: 143-156.
- FALCONER D.S. and MACKAY T.F.C. Introduction to Quantitative Genetics, 4<sup>th</sup> Edition. Longman Scientific and Technical, New York, 1996.
- 30 WELLER, J.I. *et al.* Power of daughter and granddaughter designs for determining linkage between marker loci and quantitative trait loci in dairy cattle. *J. Dairy Sci.* **73**, 2525-2537, (1990).

LYNCH M and WALSH B (1997). Genetics and analysis of quantitative traits. Sinauer Associates, Inc. Sunderland, Massachusetts.

SAS Institute Inc. 1985 SAS Users Guide: Statistics, Version 5 edition, Cary, N.C. 956p.

- 5 BOLDMAN K.G.; KRIESE L.A., VAN VLECK L.D.; VAN TASSELL C.P.; KACHMAN S.D. A manual for the use of MTDFREML, 1997.  
<http://aiipl.arsusda.gov/curtvt/mtdfreml.html>.

## SEQUENCE LISTING

5 <110> REID, SUZANNE J  
FORD, CHRISTINE A  
GEORGES, MICHEL A J  
10 COPPIETERS, WOUTER H R  
GRISART, BERNARD M J J  
15 SNELL, RUSSELL G  
SPELMAN, RICHARD J  
20  
<120> MARKER ASSISTED SELECTION OF BOVINE FOR IMPROVED MILK COMPOSITION  
25 <130> P452508 CJE  
30 <160> 21  
<170> PatentIn version 3.1  
35  
<210> 1  
<211> 11771  
40 <212> DNA  
<213> Bos taurus  
45  
<220>  
<221> CDS  
50 <222> (1)..(191)  
<223> Exon 1 CDS, determined by alignment with an amino acid sequence deduced from the cDNA  
55  
<220>  
60 <221> misc\_signal  
<222> (1)..(4)  
65 <223> these bases correspond to bases 4 to 7 of the Kozak recognition sequence. See DGAT1 cDNA for the complete recognition sequence.



<220>  
5 <221> CDS  
<222> (3809)..(3896)  
10 <223> Exon 2, determined by alignment with an amino acid sequence deduced from the cDNA

<220>  
15 <221> CDS  
<222> (5840)..(5880)  
20 <223> Exon 3, determined by alignment with an amino acid sequence deduced from the cDNA

<220>  
25 <221> CDS  
<222> (5960)..(6045)  
30 <223> Exon 4, determined by alignment with an amino acid sequence deduced from the cDNA

<220>  
35 <221> CDS  
40 <222> (6138)..(6190)  
<223> Exon 5, determined by alignment with an amino acid sequence deduced from the cDNA

<220>  
45 <221> CDS  
50 <222> (6406)..(6511)  
<223> Exon 6, determined by alignment with an amino acid sequence deduced from the cDNA  
55

<220>  
60 <221> CDS  
<222> (6601)..(6714)  
65 <223> Exon 7, determined by alignment with an amino acid sequence deduced from the cDNA

<220>  
5 <221> CDS  
    <222> (6815)..(6889)  
10 <223> Exon 8, determined by alignment with an amino acid sequence deduced from the cDNA

<220>  
15 <221> variation  
    <222> (6824)..(6889)  
20 <223> this sequence is deleted from the alternately spliced transcript.

<220>  
25 <221> CDS  
    <222> (6960)..(7063)  
30 <223> Exon 9, determined by alignment with an amino acid sequence deduced from the cDNA

<220>  
35 <221> CDS  
    <222> (7154)..(7192)  
40 <223> Exon 10, determined by alignment with an amino acid sequence deduced from the cDNA

<220>  
45 <221> CDS  
50 <222> (7271)..(7312)  
    <223> Exon 11, determined by alignment with an amino acid sequence deduced from the cDNA

55  
    <220>  
60 <221> CDS  
    <222> (7386)..(7430)  
    <223> Exon 12, determined by alignment with an amino acid sequence deduced from the cDNA  
65

<220>  
<221> CDS  
5 <222> (7505)..(7617)  
<223> Exon 13, determined by alignment with an amino acid sequence deduced from the cDNA  
10  
<220>  
<221> CDS  
15 <222> (7705)..(7770)  
<223> Exon 14, determined by alignment with an amino acid sequence deduced from the cDNA  
20  
<220>  
25 <221> CDS  
<222> (7858)..(7945)  
<223> Exon 15, determined by alignment with an amino acid sequence deduced from the cDNA  
30  
<220>  
35 <221> CDS  
<222> (8027)..(8089)  
40 <223> Exon 16, determined by alignment with an amino acid sequence deduced from the cDNA  
45  
<220>  
<221> CDS  
<222> (8162)..(8314)  
50 <223> Exon 17 CDS, determined by alignment with an amino acid sequence deduced from the cDNA  
55  
<220>  
<221> misc\_feature  
60 <222> (8315)..(8317)  
<223> translation stop codon  
65  
<220>

<221> polyA\_site  
<222> (8572)..(8578)  
5 <223>  
  
10 <220>  
<221> variation  
<222> (626)..(626)  
15 <223> thymidine (T) to guanine (G) substitution polymorphism  
  
20 <220>  
<221> variation  
<222> (3512)..(3512)  
25 <223> thymidine (T) to guanine (G) substitution polymorphism  
  
30 <220>  
<221> variation  
<222> (4040)..(4040)  
35 <223> thymidine (T) to cytosine (G) substitution polymorphism  
  
40 <220>  
<221> variation  
<222> (4963)..(4963)  
45 <223> adenine (A) to guanine (G) substitution polymorphism  
  
50 <220>  
<221> variation  
<222> (5003)..(5003)  
55 <223> guanine (G) to adenine (A) substitution polymorphism  
  
60 <220>  
<221> variation  
<222> (5997)..(5997)  
65 <223> cytosine (C) to thymine (T) substitution polymorphism

- <220>
- 5 <221> variation
- <222> (6829)..(6830)
- 10 <223> adenine (A)-adenine (A) to guanine(G)-cytosine (C) substitution p  
olymorphism  
AA corresponds to the Q allele  
GC corresponds to the q allele
- 15 <220>
- <221> variation
- 20 <222> (6892)..(6892)
- <223> guanine (G) to adenine (A) substitution polymorphism
- 25 <220>
- <221> variation
- 30 <222> (7224)..(7225)
- <223> guanine (G)-guanine (G) to adenine (A)-cytosine (C) substitution  
polymorphism  
GG-AC
- 35 <220>
- 40 <221> variation
- <222> (7438)..(7438)
- <223> adenine (A) to guanine (G) substitution polymorphism
- 45 <220>
- 50 <221> variation
- <222> (7456)..(7456)
- <223> cytosine (C) to thymidine (T) substitution polymorphism
- 55 <220>
- 60 <221> variation
- <222> (7987)..(7987)
- <223> guanine (G) to adenine (A) substitution polymorphism
- 65

<220>  
<221> variation  
5 <222> (8402)..(8402)  
<223> ctosine (C) to thymidine (T) substitution polymorphism  
10 <220>  
<221> misc\_feature  
15 <222> (9434)..(9434)  
<223> ambiguous nucleotide  
20 <220>  
<221> misc\_feature  
25 <222> (9496)..(9496)  
<223> ambiguous nucleotide  
30 <220>  
<221> misc\_feature  
35 <222> (10402)..(10417)  
<223> ambiguous nucleotides  
40 <220>  
<221> primer\_bind  
45 <222> (6579)..(6601)  
<223> Primer 17F  
CCTGAGCTTGCCTCTCCCACAGT  
50 <220>  
<221> primer\_bind  
55 <222> (7036)..(7058)  
<223> Primer 18R  
CCAGGAGTCGCCGCAGCAGGAAG  
60 reverse primer  
65 <220>  
<221> primer\_bind

5 <222> (7280)..(7303)  
<223> Primer 6F  
CCGGCCATCCAGAACTCCATGAAG

10 <220>  
<221> primer\_bind  
<222> (7585)..(7605)  
<223> Primer AW446985 dn1  
15 TAGAACTCGCGGTCTCCAAAC  
reverse primer

20 <220>  
<221> primer\_bind  
<222> (8222)..(8244)  
25 <223> Primer InsUp1  
TGGCTGTCACTCATCATCGGGCA

30 <220>  
<221> primer\_bind  
35 <222> (8566)..(8589)  
<223> Primer 14R2  
TTGCACAGCACTTTATTGACACA

40 <220>  
<221> primer\_bind  
45 <222> (6813)..(6830)  
<223> Primer SNP1\_FAM  
50 AGC TTT GGC AGG TAA GGC

<220>  
55 <221> primer\_bind  
<222> (6813)..(6830)  
<223> Primer SNP1\_HEX  
60 AGC TTT GGC AGG TAA GAA

<220>  
65 <221> primer\_bind

5 <222> (6831)..(6844)  
<223> Primer SNP1\_2P  
GGC CAA CGG GGG AG

<220>

10 <221> primer\_bind  
<222> (7424)..(7438)  
<223> Primer SNP2\_FAM  
15 GCT GGC GGT GAG TGA

<220>

20 <221> primer\_bind  
<222> (7424)..(7438)  
<223> Primer SNP2\_HEX  
25 GCT GGC GGT GAG TGG

<220>

30 <221> primer\_bind  
<222> (7439)..(7453)  
<223> Primer SNP2\_2P  
35 CCT GCT GGG TGG GGA

<220>

40 <221> primer\_bind  
<222> (7442)..(7456)  
<223> Primer SNP3\_FAM  
45 GCT GGG TGG GGA CGC

<220>

50 <221> primer\_bind  
<222> (7442)..(7456)  
<223> Primer SNP3\_HEX  
55 GCT GGG TGG GGA CGT

<220>

60 <221> primer\_bind



5 <222> (7457)..(7470)  
<223> Primer SNP3\_P  
GTG GGG GCG GGT GG

10 <220>  
<221> primer\_bind  
<222> (8388)..(8402)  
<223> Primer SNP4\_FAM  
15 TGC CCC AAC CTG GGT

20 <220>  
<221> primer\_bind  
<222> (8388)..(8402)  
<223> Primer SNP4\_HEX  
25 TGC CCC AAC CTG GGC

30 <220>  
<221> primer\_bind  
<222> (8403)..(8417)  
<223> Primer SNP4\_2P  
35 GCA GCA GGA GGA GGC

40 <220>  
<221> primer\_bind  
<222> (6811)..(6830)  
<223> Primer Dgat 21  
GTAGCTTTGGCAGGTAAGAA

50 <220>  
<221> primer\_bind  
<222> (6965)..(6984)  
<223> Primer Dgat 22  
GGGGCGAAGAGGAAGTAGTA  
60 reverse primer

65 <220>  
<221> primer\_bind

5 <222> (6613)..(6632)  
<223> Primer Dgat 23  
TGGCCCTGATGGTCTACACC

10 <220>  
<221> primer\_bind  
<222> (6829)..(6850)  
<223> Primer Dgat 24B  
15 GGGCAGCTCCCCCGTTGGCCGC  
reverse primer

20 <220>  
<221> primer\_bind  
<222> (6651)..(6670)  
25 <223> Primer DgatforAD  
TTCTCCTACCGGGACGTCAA

30 <220>  
<221> primer\_bind  
35 <222> (6871)..(6890)  
<223> Primer ReverseNZ  
CCGCGGTAGGTCAGGTGTC  
40 reverse primer

45 <220>  
<221> primer\_bind  
<222> (6825)..(6838)  
50 <223> Probe ForAA (FAM)  
CGTTGGCCTTCTTA

55 <220>  
<221> primer\_bind  
60 <222> (6823)..(6836)  
<223> Probe DgatADGC (VIC)  
TTGGCCGCCTTACC  
65

<220>  
<221> primer\_bind  
5 <222> (6651)..(6670)  
<223> Primer DgatforAD  
TTCTCCTACCGGGACGTCAA  
10  
<220>  
15 <221> primer\_bind  
<222> (6878)..(6972)  
<223> Primer DgatrevAD  
20 AAGTAGTAGAGATCGCGGTAGGTCA  
reverse primer  
<220>  
25 <221> primer\_bind  
<222> (6825)..(6838)  
30 <223> Probe ForAA (FAM)  
CGTTGGCCTTCTTA  
35  
<220>  
40 <221> primer\_bind  
<222> (6823)..(6836)  
<223> Probe DgatADGC (VIC)  
45 TTGGCCGCCTTACC  
<220>  
50 <221> primer\_bind  
<222> (6652)..(6671)  
55 <223> Primer DgatforRT66  
TCTCCTACCGGGACGTCAAC  
60  
<220>  
<221> primer\_bind  
65 <222> (6874)..(6964)  
<223> Primer DgatrevRT66

GAGATCGCGGTAGGTCAGGTT  
reverse primer

5  
<220>  
<221> primer\_bind

10 <222> (6711)..(6972)  
<223> Primer DgatforRTless66, GCTGCTTTGGCAGATCTCTACTACTT  
This primer was designed to selectively bind and amplify the cDNA  
splice variant. The corresponding binding site in this genomic s  
equence comprises bases 6711 to 6715, 6815 to 6823 and 6960 to 69  
72.

20 <220>  
<221> primer\_bind  
<222> (7022)..(7038)

25 <223> Primer DgatrevRTless66  
AAGCGCTTTCGGATGCG  
reverse primer

30  
<220>  
<221> primer\_bind

35 <222> (6857)..(6870)  
<223> Probe Dgatwith66 (FAM)  
CCGTGAGCTACCC

40  
<220>  
<221> primer\_bind  
<222> (6976)..(6990)

50 <223> Probe Dgatless66 (VIC)  
CTTCGCCCCCACCCT

55  
<400> 1  
atg ggc gac cgc ggc ggc gcg ggc ggc tcc cgg cgc cgg agg acg ggg 48  
Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Arg Thr Gly  
1 5 10 15  
60 tcg cgg cct tcg atc cag ggc ggc agt ggg ccc gcg gca gcg gaa gag 96  
Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu  
20 25 30  
65 gag gtg cgg gat gtg ggc gcc gga ggg gac gcg ccg gtc cgg gac aca 144  
Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr  
35 40 45

	gac aag gac gga gac gta gac gtg ggc agc ggc cac tgg gac ctg ag	191
	Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asp Leu Arg	
	50 55 60	
5	gtagcgggtgc gcgtgacccc taacctttga cccctgatac ggggcccctg cgacccaacc	251
	tggtggccca ggctgtcgg cggcagctcg ggctcgagtc cgagagtctg gcgcctggac	311
10	cttgggtcac agctgtgccc ctccggcctc cacggggaaa cttagcgga ggttgggggc	371
	ggaggggtctc ctgcccggaa cccccaggta cggggggcca ggggaggga gcggctcaac	431
15	ttctagacgc cctccctctg ccttcctttg gtgggttctg aagctttccc agggtagacc	491
	cactacgcac agtgtcctct acctggaagg agatacagg gtccttctctg agggctatga	551
	gggggtgcctt gtgggttgat aaagctccc ggggaggagg gtggaccggc ggagaacaga	611
20	ggcaggggca gtgctagggg atttctcatc cctcgagac cctccagaga atggtcttca	671
	caaaggtccc tcacccgtca cccggcgatt gactggccta ggatcctgct tattaccagc	731
25	acaaatggct gctctagggg caaagtgggt cctgtaatgg gaccctcacc cctggttggg	791
	gtacagggga ggagttggaa gtgcgcacac ccacaggtgg gcgcctctgct tagctgaagg	851
	actgatggga aggagttggg ggagcaagct gcggctgaaa gggaggatct gaccacgtg	911
30	ggcatcagct aagtcctgct ggctgcctcc aggcgttccc tttgccatcc tccacgcccc	971
	tccccgggg cctgaccttc atcctgggtca agggctctca ggggctctgg ttttgggac	1031
35	agctccagag cttagaggtta tcaaggagga agtgggcaac aggtcagtc gcaaggattt	1091
	gctatcttca ctgggtgctg tggggagggg agggacaagg gcagttgggg tgcaggcact	1151
	gtccctgccc ttggggggca cacagttcac ctgagagata agatagccgc agccctgaag	1211
40	agttagagca aaggtcaggc acagagttca ggatgacacc aggggagggt ggctctgtga	1271
	ggggcactgg ctctctacag gccccagggt gtcctgaggg ggcggtgca aaggccagga	1331
45	ggcccacagg cccctctgcc cactcctggg gaactggatt tggggtcact ttgtatgagg	1391
	tggggcgggg taccagcttt gggccaagct gtcaccctgg atgggccatc acttgccctg	1451
	tctgtatagg ccagatggcc agaagctgct cctgtcctgt tgatggcca tctcagagg	1511
50	ctggaccctc gggaagagga gcagttggtg gcagggatgg gccaccggag accctcctga	1571
	cctccaggac acgcagctgt gtgtgcctgt cccaagcca catgccacat ggctaggggc	1631
55	ctcctggggc agggctgggc attggtcttg ctactcttgg tatcgccat ggcttccctg	1691
	cctcccagtc atcatcctcc cacctctgcc tcctgctg ttctctcttt tctcctcagg	1751
	cccttcggga catttctgc tcacctagg ctgggcaggc ggggtcagg gcccgggtgtg	1811
60	agctcactcc ttccggcagc aaggtgtagc tatgtgccg aaggaaggcc gctgctgttg	1871
	cctcgctct gagtgatcc ctccaggtc ctccacact cctgtgccc cgacacctgg	1931
65	tgcgtccttc agccattggt tcatgtgtcc tccaggcaca gctttctagt ccagagcctc	1991
	taggctgggt gcaggaagt ctgaggaagt ggcagccggg aggcgagctg gcaccctgtc	2051

61

	cctccttggt ctgtccgtcc ctgcccctgg accgtatggc cccgcatgtg tgatccccac	2111
	ttggggctgt gcctctgggc aagttaggaa gcttggtgag cctcattttc atgtgcccgc	2171
5	ctcccagtac tgatgtgcag gttgaatgag gtgccaaactg taatgagttg gaatggccct	2231
	gctggctggg tgggactggg gagcaggtgg gggccgctgg ggggcacaga ggcacacca	2291
10	gtgcctcagt cagggagagg gtgacagaga agctctgggt gagggccac ctccactctg	2351
	gccatggctg ctgccctttg gtccactgca gtgaactgtg ccatggggct ggacctctgt	2411
	ggggattggt gggcagtggt ctttcttccc gcttggggcc tctgacctct gggggcaggg	2471
15	cgctgcccgg gtgggacagt cggaaggctg gtagagggac ctgaggggtc tgtgtggtgg	2531
	ctgggggcag gcctcaggaa ttgacagca gggatctgga aaagctttaa taacattatt	2591
20	tgttgtcagg attgggaaat gctccccctc cccctcccc tctttcatct tagagactgc	2651
	tgcacatctg gtcagtgtgg tcttcttggg gggccccaag gtggcagggg tcacactgtt	2711
	atgaaaccgt cccctgggta tgtggtgcag acatgcacat gcagatgggt attggcaggt	2771
25	tgtagcatga ggtggctttg ggacggttcc agtgacagtg agtgggctgg atctgggggg	2831
	ttctgggcag gtccatcaag cggatacccc cacagactgt cctcttgga tagttgggcc	2891
30	tgggagccct gcttgccctg caaaaggca ggcgcagagt catgaagaag agggcttggg	2951
	ggctcagagc cccactgtgt gtgcagccca gggtagacct ggaggagggt cgtgggcagg	3011
	ctgggccggc cggggcgggg ggtggggggg cctggtgtga aagggaccca gggccagact	3071
35	gtcagcgtg cctggctgag gatgctggca ccctgtcctc cccagccgtc tgtctcctgg	3131
	gtgcagccat ctgagtgtg accccagccg cccctggagg ctggctgttc tctgtgccc	3191
40	tattgctggg gacatgtgtc cacaggaggg aaagggaagc cccggcctct ccccttaca	3251
	aactggaggc cttgtcfaat gccctggatg gcctcctggt ggcaggggtg ttggtgggag	3311
	gtggggctgc tgcttagaac ccgccagcg gcctgggcct gggctgagct gcacccctcc	3371
45	acctctgct ccagctgagg gttggttcc atctccacca ggcagcgac tgggcacagg	3431
	gctctcagag gcaggtctg aaagtcccct gctggcttct gcagtgact ccaggcgccg	3491
50	agcccccagg gggctcgc atgcgtcacc ctgcgaagcc acgtgaaggc tgggtcctcc	3551
	cctccggaag ggccaaatgc agggcatggg tggtttgaat ggtggccct gggctccccg	3611
	gagggaccag ctgctgtgag ggccgcccc tccccacttc cgtcttgcat caccagctcc	3671
55	tgtggcactc cccacgcccc gtccccagt gggagcgga gggcccggt ggctctgccc	3731
	gcggaggggg atgtgtggg ggcggggtg ccttgctgcc agatgctctg ccccgagtgt	3791
60	ccgtctccgc tctccag g tgt cac cgc ctg cag gat tcc ctg ttc agt tct	3842
	Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser	
	65 70 75	
65	gac agt ggc ttc agc aac tac cgt ggc atc ctg aat tgg tgt gtg gtg	3890
	Asp Ser Gly Phe Ser Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val	
	80 85 90	
	atg ctg gtacgtagag tgacacctg gagcaagggt cctgacggcc ggggggcat	3946

Met Leu

5	gggctcttct ccaggggtag gtgtctgtac ttgtgtagct gtggtgaatg gagctctgtg	4006
	ctggcgggtgg ggggtccctgg agcagccgta ccctgggacc ctaccgggag catgctcatg	4066
	ccgtccctgc tgaatcccag gagatgctg cagagggcag cctgggagcc tctgagctgg	4126
10	ggtctgcgcc ccagggggca ctggagtctc ccagggggc gagagagagt aggcagggat	4186
	ggtctggtgg ccctgggtgg gggatggctg ctccgtgggc ccaggccctc cctggcagca	4246
15	caggtagtg gtcttggggg tccacgtaga acttctctt ctgttccaa ttgccctcat	4306
	gggtgcggca tgcttgggtg aacctggggg agcaggggta ggacatgctt ctacagcccag	4366
	cccacagctc caggccacac tctgcaggac tctggcccct ccctcagccc tggaggggagc	4426
20	aggactggag tcctgtgtcc gccttgtctt gacctggccg aggccactgc tgtggggccc	4486
	cagcaggcct gccacagaga aggtggagt cagggacccc aggggcagcc ttcagggtgg	4546
25	ggcaggggta ggcccagctg ggcccagccc caccgctcag tgctgatgtg gcgcgaggcc	4606
	ttcggccctc cagctgacgt gtctgcctgc cctgggtgtg gctccagagg ctgcctgtgt	4666
	accaggggcc cccacgcttc tgtttgtggt tctgggcagt cccctgggga gcggtggggg	4726
30	ctgtgtgcca gtccagacct agtagtcac gcgtcctggt ctctggaggc cgtggctggt	4786
	ccaggactgt ggcaagggtg tcgtgcaggg caggccctca gcagcctgtc tgttctcctg	4846
35	cagccccag cctcctggcc ctttgggtgca cccacaaagc tccccctcc cccaggagct	4906
	ggggccgctt gctgcgtcct ctccgcagcc tgggcttcca ggtggctggg cctcttagca	4966
	gctccaaact ttgcctgtgg tgggctctca ggacaggcaa ctgccagtcg gcagacattg	5026
40	caggaccacg tgtgtcctgg taagctggct ggttaggtgt ttagctgggg gatggtgtgg	5086
	cagggtggcc ctgcctctct gagcctgtca cctcctcggg aagccttctg ggtgggggac	5146
45	tccacccatg tcgcctggag aagcatcact tttccacaga gccttctgca acccccgtgg	5206
	ggcctgagcc tgggggtggg gaggtggtgg cccctgctcc tgcagaggcc agccaggcat	5266
	ctggccccag gccactggca agagctcgtt gtgttggggg atctgtcctt tgcctgtgct	5326
50	gcaggagcgg ccgaggcagg cgggggcgtg agtaggggtg gagaccagc cccagcttcc	5386
	ccagccctc aggaccggcc tgctctttcc caccaccca ccaagtgcgt gggcacaccc	5446
55	cgctgtgag gatgggccc gttggcaggg cggagccctg ggagggtggc agtgcgccg	5506
	gcaggcttgg acttactgg ggttgggggt tgcgtgtgtg gccaggggag ctgaccgct	5566
	tgggtggagc gacggccgct gggcagcagg tttcttctgc cacgggtggca caggcacctg	5626
60	gggttgtggt tggctccagg cgggcggggg ctgcgtgccc ctgcgcaggc acataggccg	5686
	tgggtgggga gtctcagagc ttggcgtgag gtcccacagg gctgggcctg caggatggag	5746
	gccactgtcc tgagctgcag gtgctggcag gagctggggg gggcgttctg gggccgtggc	5806
65	tgacagcgtt atgtccctct ctctctatcg cag atc tta agc aac gca cgg tta	5860
	Ile Leu Ser Asn Ala Arg Leu	

		63	
		95	100
5	ttt cta gag aac ctc atc aa gtgagtgggc cccggcctgc cccagcccct Phe Leu Glu Asn Leu Ile Lys 105	5910	
	gccacctcac cctcgccta cacagaccct caccacctg cgtctgcag g tat ggc Tyr Gly	5966	
10	atc ctg gtg gac ccc atc cag gtg gtg tct ctg ttc ctg aag gac ccc Ile Leu Val Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro 110 115 120 125	6014	
15	tac agc tgg cca gct ctg tgc ctg gtc att g gtgagctggg tgcccaggag Tyr Ser Trp Pro Ala Leu Cys Leu Val Ile 130 135	6065	
20	gcctcaggcc ggcgtgggt gggacagggc tgatctgggc ctgaacctgc cctgggttgc ttctgtcctc ag tg gcc aat atc ttt gcc gtg gct gcg ttc cag gtg gag Val Ala Asn Ile Phe Ala Val Ala Ala Phe Gln Val Glu 140 145	6125 6175	
25	aag cgc ctg gcc gtg gtaagcagt ccctcagcc ctcccctgac ttgcctcaag Lys Arg Leu Ala Val 150	6230	
30	gtccttacca gtcgggctta gggcgggcca ccagctggtc cactgtgtct tcagggtttt gggcctttcg tggccttctc gagaggggct gcacctcagg cctgggtggct cttcctcagg gaggtcctct gaccaggag gggggtccct ggctgacgct ctgctccac cccag gga Gly	6290 6350 6408	
35			
40	gct ctg acg gag cag gcg ggg ctg ctg ctg cac ggg gtc aac ctg gcc Ala Leu Thr Glu Gln Ala Gly Leu Leu Leu His Gly Val Asn Leu Ala 155 160 165 170	6456	
	acc att ctc tgc ttc cca gcg gcc gtg gcc ttt ctc ctc gag tct atc Thr Ile Leu Cys Phe Pro Ala Ala Val Ala Phe Leu Leu Glu Ser Ile 175 180 185	6504	
45	act cca g gtgggcccc ccccccccc cgccccgcc cacgtgtct cggccacggg Thr Pro	6561	
50	cagcgcgggg ggcgtggcct gagcttgcct ctcccacag tg ggc tcc gtg ctg Val Gly Ser Val Leu 190	6614	
55	gcc ctg atg gtc tac acc atc ctc ttc ctc aag ctg ttc tcc tac cgg Ala Leu Met Val Tyr Thr Ile Leu Phe Leu Lys Leu Phe Ser Tyr Arg 195 200 205	6662	
60	gac gtc aac ctc tgg tgc cga gag cgc agg gct ggg gcc aag gcc aag Asp Val Asn Leu Trp Cys Arg Glu Arg Arg Ala Gly Ala Lys Ala Lys 210 215 220 225	6710	
	gct g gtgagggctg cctcgggctg gggccactgg gctgccactt gcctcgggac Ala	6764	
65	cggcaggggc tcggctcacc cccgacccgc ccctgccgc ttgctcgtag ct ttg Ala Leu	6819	



5	gca ggt aag aag gcc aac ggg gga gct gcc cag cgc acc gtg agc tac Ala Gly Lys Lys Ala Asn Gly Ala Ala Gln Arg Thr Val Ser Tyr 230 235 240	6867
	ccc gac aac ctg acc tac cgc g gtgaggatcc tgccgggggc tggggggact Pro Asp Asn Leu Thr Tyr Arg 245 250	6919
10	gcccggcggc ctggcctgct agccccgcc tcccttcag at ctc tac tac ttc Asp Leu Tyr Tyr Phe 255	6973
15	ctc ttc gcc ccc acc ctg tgc tac gag ctc aac ttc ccc cgc tcc ccc Leu Phe Ala Pro Thr Leu Cys Tyr Glu Leu Asn Phe Pro Arg Ser Pro 260 265 270	7021
20	cgc atc cga aag cgc ttc ctg ctg cgg cga ctc ctg gag atg Arg Ile Arg Lys Arg Phe Leu Leu Arg Arg Leu Leu Glu Met 275 280 285	7063
25	gtgaggcggg gcctcgcggg ccagggtggg cgggcctgcc ggcacccggc accggggctc	7123
	agctcactgt ccgcttgctt ccttccccag ctg ttc ctc acc cag ctc cag gtg Leu Phe Leu Thr Gln Leu Gln Val 290	7177
30	ggg ctg atc cag cag gtacgtgcc gggggggggg gggggggggg gggggggact Gly Leu Ile Gln Gln 295	7232
35	ctggggccgt tggggagctg actctgcgct ttttgtag tgg atg gtc ccg gcc atc Trp Met Val Pro Ala Ile 300 305	7288
40	cag aac tcc atg aag ccc ttc aag gtgagcaggc aggcctggca ggggtgggttc Gln Asn Ser Met Lys Pro Phe Lys 310	7342
	cggggtcagg gctgaggag ccagctgtgc cctgtgccca cag gac atg gac tac Asp Met Asp Tyr 315	7397
45	tcc cgc atc gtg gag cgc ctc ctg aag ctg gcg gtgagtgacc tgctgggtgg Ser Arg Ile Val Glu Arg Leu Leu Lys Leu Ala 320 325	7450
50	ggacgcgtgg gggcgggtgg ggctgttctg gcacctggca cccactcccc acag gtc Val	7507
55	ccc aac cac ctc atc tgg ctc atc ttc ttc tac tgg ctc ttc cac tcc Pro Asn His Leu Ile Trp Leu Ile Phe Phe Tyr Trp Leu Phe His Ser 330 335 340 345	7555
60	tgc ctg aac gcc gtg gct gag ctc atg cag ttt gga gac cgc gag ttc Cys Leu Asn Ala Val Ala Glu Leu Met Gln Phe Gly Asp Arg Glu Phe 350 355 360	7603
	tac cgg gac tgg tg gtgggtggcc ttgccccggc gggggtggtg ggggccccg Tyr Arg Asp Trp Trp 365	7657
65	ctggggctgg ggccggagcc cctgcccact ctgccccgcc cccgcag g aac tcc gag Asn Ser Glu	7714

65

	tcc atc acc tac ttc tgg cag aac tgg aac atc cct gtt cac aag tgg	7762
	Ser Ile Thr Tyr Phe Trp Gln Asn Trp Asn Ile Pro Val His Lys Trp	
	370 375 380 385	
5	tgc atc ag gtgggtgtgc gcctgggggc ggggggttgg ggggtgggac	7810
	Cys Ile Arg	
10	gggggtcgcgt ggccccggcg cccagcccac tgccgcctcc cccgcag a cac ttc tac	7867
	His Phe Tyr	
	390	
	aag ccc atg ctc cgg cgg ggc agc agc aag tgg gca gcc agg acg gca	7915
15	Lys Pro Met Leu Arg Arg Gly Ser Ser Lys Trp Ala Ala Arg Thr Ala	
	395 400 405	
	gtg ttt ctg gcc tcc gcc ttc ttc cac gag gtcagtgcac tgagggcgcg	7965
	Val Phe Leu Ala Ser Ala Phe Phe His Glu	
20	410 415	
	ccctgcccct ggtgggggtg ggggtggggg tgggggctcg ctgacgcccc tctcccctca	8025
	g tac ctg gtg agc atc ccc ctg cgc atg ttc cgc ctc tgg gcc ttc acc	8074
25	Tyr Leu Val Ser Ile Pro Leu Arg Met Phe Arg Leu Trp Ala Phe Thr	
	420 425 430	
	ggc atg atg gcg cag gtgagcagcc ctggaccccc gctccgcccc gccccgcgag	8129
	Gly Met Met Ala Gln	
30	435	
	cgcagaggct cactcccgtc ctgtgtcccc ag atc ccg ctg gcc tgg ata gtg	8182
	Ile Pro Leu Ala Trp Ile Val	
	440 445	
35	ggc cgc ttc ttc cgc ggc aac tac ggc aac gcg gcc gtg tgg ctg tca	8230
	Gly Arg Phe Phe Arg Gly Asn Tyr Gly Asn Ala Ala Val Trp Leu Ser	
	450 455 460	
	ctc atc atc ggg cag ccg gtg gcc gtc ctg atg tac gtc cac gac tac	8278
40	Leu Ile Ile Gly Gln Pro Val Ala Val Leu Met Tyr Val His Asp Tyr	
	465 470 475	
	tac gtg ctc aac cgt gag gcg ccg gca gcc ggc acc tgagcgctc	8324
45	Tyr Val Leu Asn Arg Glu Ala Pro Ala Ala Gly Thr	
	480 485	
	caggctggcc ccctcgtggg tggttgactg ctttgccgcg ctgcctgcgg ctggactaga	8384
50	gcctgcccga acctggggcg agcaggagga ggctgtgctg gtggaagctg cctcctggcc	8444
	tccaccaggc ctctgcctga agggcttcct cctgccaggg gagagcaggc ccgacgcagt	8504
	tctggcccct gggaggtgcc catgctctgg aaaccctaca gatctcgccc aagggtctga	8564
55	atgtgtcaat aaagtgtgtg gcacagttag ctccctcagc ctccagggca cagggtgtggc	8624
	aggagggggc ggccctccca cgtggggcca tgctgtggga aggaggcccc agcgcttgga	8684
60	gaggagctgg ggctgtggtg accctccctg cctcacaggg ctctgtgggc agacgtcttg	8744
	ccctgcaagg tggagactcc atgctccaag gccccctgtg cctgaggtct gcacacaagt	8804
	ggattcaact tgggtcaggc cagaggctaa ggtgtggaag aagggtgaga atcaggctga	8864
65	cttgaacggc agcaaagact ccaaggcaag gctgcagaag tctcaaaagc tatgcgcaca	8924
	gtcccctgct ggggtgctca cctgggctgg gctctgggct gcttgacaaa agcaggtggc	8984

	ctggctcagc cctcaccgag ggccttcctt gggggcagag gttggcctga tgccaggggc	9044
5	tccccgtttt tccaggccct cagcaggtag ttgggtgtgg ccctcatgat accttggctc	9104
	cagagctccg ccaactcaaaa agcttggcag tgaggcaagg gcaaccccg gctgttcccc	9164
	cctctactgg ctctgccgcc tgggttgga accctgaggc tgtgccaggc aggtgtaccc	9224
10	tgacagccag ccatggccca gtaagatggg tgcccagagt ggtacctggg cagcggaccc	9284
	agctgtgctg cccccgccc aaccagaagc cgctctagcc catggtggtc gtctgggcca	9344
15	gacaggctgg ttggctaggc actgtttgtt ctacagcagg ttaggcagc gtctccctga	9404
	cccctgcctt ctagggaagc accaccctgn gccctactca tcagcaagga cagcgagcag	9464
	ggctgagctg ggggtgcgtg ggctgctacg gnccccgcac cttcatcaca tgcacctctg	9524
20	cacccccctg tgccctgactc aagagtgggg gggggggtcc tgtgcttctt tcattccaga	9584
	cccacggtgc tgaccagtg caccacctg gtccgctagt gctgacctgg ccacagggct	9644
25	cctgtgggcc cagctgata ccgcccgtt cccttcataa agaactcttg agcacatgca	9704
	gcccagggga gccaggaggc tccagtgtgc tgtgtccatc tgccctccctt cagcccttc	9764
	cgagacactg cgcacatgc cccctccac ccccaaccac actggcagga ggaacacaca	9824
30	gggagaccac acacagagct cgttgtttat aaatctctgc ctggctcatc ggtctgtttg	9884
	tccatgtata tatctgcata tctctatgga aggggaaagg gggactcgtg taaaaatcca	9944
35	aaatacaatt ctatgaacac ctgcatcctg gtcagtctga gtgtggccgt gaagcccagg	10004
	tgagctgtgg ctacagggc taggcctcgt gtgctggccg ggggccactc cccaccccct	10064
	ctccccccct ccgccagcca ggggaccagg ctctctggaca ccaggcctgc ccaaggcctg	10124
40	ctctctctct ggggcttcta cgagacagtg gggctccttg ctttgggggg ttctgagccc	10184
	gtcagcaggg agatggtggg gtcacatctt atatttcgtc tccctcgag aagtaggagc	10244
45	cctccccag ctggaagagc accggcaggt cgctgctccc cacgtccacg gagccgggt	10304
	ccaggagcag caggggctgg gcggtgtagt gcaccagctg cttccctagg ggtgcgactg	10364
	ggtcagggtg ccggtggggc cggggggcgg ggtggggnnn nnnnnnnnnn nnncccccc	10424
50	ccccggcccc agccacccc cctacgcacg ctggccaggc tgctgtccag gtcgggcagg	10484
	ctcatgtcgg gcaccgtaac cgaggggctg aacagctgca ggaagaggg gcgggtcaga	10544
55	ctgccctgga agcggggcgg gctgggcggg gggcgggcag ggttagactc acatccagca	10604
	gggtgctggt gtccacgctg aagccatggc ttgtcagcat ggtctgcagg ttgtccaggt	10664
	tggagtccat ggcgtccaag tggctgctga gctcggctct gccagagga aggggagcag	10724
60	gtgacgtggc atccaggcac cccagggtgc agtctgccg gcccttaagc ccagctgcca	10784
	gcctgccctg cgcaggtca cgggaggccc tgaggtgggg ccctgggcct gggacacttg	10844
65	gccaccggtg gtggctcagg ctccctcatc accgtgaggc cccgtaaggc agcacgtcag	10904
	gctgggccgg ctaagacatc agggggctga gttcagggtc cagaagggcc cggggcccca	10964

67

gcttctcagg ttcagcaacg cctgcccttc acactggagg acagcaggag ggtggagctg 11024  
tgctcggcca ggctgggcca gcctacccca gcatgcgcca ggatagatta ggcctgcccc 11084  
5 ctcgaggagg ccaggcagta cctcggggtg ctgcctcagg cccagaggt ggtggggag 11144  
cctgttcaca ggctgggcag ggtctgctc agaggctggg gaggggcccc agctctggcg 11204  
gagccctcct ccctccccag tgggacagcg ctaaccctgg ctggactcgg ccatgcagag 11264  
10 aggaggaggg gcaggggaag aggcgggcga cccagacat ctttgagtg cgagccaaac 11324  
tgcaagatag aaagacaaga gccccctct gcgggtgggg cccagcgagt ccaagcccc 11384  
15 caccctgttc acccacctct ggggtggtag gacagacaga ccaggcgcc cgaggagggc 11444  
ggctcagggg gcaggggtgc ccgccacgtg cctggttccc acggtgcagg ggctgcggga 11504  
ccccctggct ggcatgtctc cctccccagc tccatgtctt ggccagggtc ggccagtttc 11564  
20 cccagcaaa tcccatgccg agggcctgag ggccatgcgt gtccagccc gactgtcct 11624  
tgtcggaggc ccctgcgctt ccagagaccg aagggcgctt ggaaggcact cacttgtcta 11684  
25 ggcaggcgac gctgaggcac ttctcgggag ccgaggcggg gagggtcgac ggatagcggc 11744  
cccgagtgat ccgatagaag cttcgta 11771

30 <210> 2  
<211> 489  
<212> PRT  
35 <213> Bos taurus

40 <220>  
<221> misc\_feature  
<222> (8315)..(8317)  
45 <223> translation stop codon  
<220>  
50 <221> misc\_feature  
<222> (9434)..(9434)  
<223> ambiguous nucleotide  
55 <220>  
<221> misc\_feature  
60 <222> (9496)..(9496)  
<223> ambiguous nucleotide  
<220>  
65 <221> misc\_feature

68

&lt;222&gt; (10402)..(10417)

&lt;223&gt; ambiguous nucleotides

5 &lt;400&gt; 2

Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Thr Gly  
 1 5 10 15

10 Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu  
 20 25 30

15 Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr  
 35 40 45

20 Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asp Leu Arg  
 50 55 60

25 Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser Asp Ser Gly Phe Ser  
 65 70 75 80

Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val Met Leu Ile Leu Ser  
 85 90 95

30 Asn Ala Arg Leu Phe Leu Glu Asn Leu Ile Lys Tyr Gly Ile Leu Val  
 100 105 110

35 Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro Tyr Ser Trp  
 115 120 125

40 Pro Ala Leu Cys Leu Val Ile Val Ala Asn Ile Phe Ala Val Ala Ala  
 130 135 140

45 Phe Gln Val Glu Lys Arg Leu Ala Val Gly Ala Leu Thr Glu Gln Ala  
 145 150 155 160

Gly Leu Leu Leu His Gly Val Asn Leu Ala Thr Ile Leu Cys Phe Pro  
 165 170 175

50 Ala Ala Val Ala Phe Leu Leu Glu Ser Ile Thr Pro Val Gly Ser Val  
 180 185 190

55 Leu Ala Leu Met Val Tyr Thr Ile Leu Phe Leu Lys Leu Phe Ser Tyr  
 195 200 205

60 Arg Asp Val Asn Leu Trp Cys Arg Glu Arg Arg Ala Gly Ala Lys Ala  
 210 215 220

65 Lys Ala Ala Leu Ala Gly Lys Lys Ala Asn Gly Gly Ala Ala Gln Arg  
 225 230 235 240

Thr Val Ser Tyr Pro Asp Asn Leu Thr Tyr Arg Asp Leu Tyr Tyr Phe

69

	245	250	255
5	Leu Phe Ala Pro Thr Leu Cys Tyr Glu Leu Asn Phe Pro Arg Ser Pro 260 265 270		
10	Arg Ile Arg Lys Arg Phe Leu Leu Arg Arg Leu Leu Glu Met Leu Phe 275 280 285		
15	Leu Thr Gln Leu Gln Val Gly Leu Ile Gln Gln Trp Met Val Pro Ala 290 295 300		
20	Ile Gln Asn Ser Met Lys Pro Phe Lys Asp Met Asp Tyr Ser Arg Ile 305 310 315 320		
25	Val Glu Arg Leu Leu Lys Leu Ala Val Pro Asn His Leu Ile Trp Leu 325 330 335		
30	Ile Phe Phe Tyr Trp Leu Phe His Ser Cys Leu Asn Ala Val Ala Glu 340 345 350		
35	Leu Met Gln Phe Gly Asp Arg Glu Phe Tyr Arg Asp Trp Trp Asn Ser 355 360 365		
40	Glu Ser Ile Thr Tyr Phe Trp Gln Asn Trp Asn Ile Pro Val His Lys 370 375 380		
45	Trp Cys Ile Arg His Phe Tyr Lys Pro Met Leu Arg Arg Gly Ser Ser 385 390 395 400		
50	Lys Trp Ala Ala Arg Thr Ala Val Phe Leu Ala Ser Ala Phe Phe His 405 410 415		
55	Glu Tyr Leu Val Ser Ile Pro Leu Arg Met Phe Arg Leu Trp Ala Phe 420 425 430		
60	Thr Gly Met Met Ala Gln Ile Pro Leu Ala Trp Ile Val Gly Arg Phe 435 440 445		
65	Phe Arg Gly Asn Tyr Gly Asn Ala Ala Val Trp Leu Ser Leu Ile Ile 450 455 460		
	Gly Gln Pro Val Ala Val Leu Met Tyr Val His Asp Tyr Tyr Val Leu 465 470 475 480		
	Asn Arg Glu Ala Pro Ala Ala Gly Thr 485		
	<210> 3		
	<211> 31		

<212> DNA  
<213> Bos taurus

5  
<220>  
<221> misc\_signal  
10 <222> (29)..(31)  
<223> bases 1 to 3 of the Kozak recognition sequence. See the genomic s  
15 equence from the start codon for bases 4 to 7 of the Kozak recogn  
ition sequence or the DGAT1 cDNA for the complete recognition seq  
uence.

20 <400> 3  
acttggccgc ggcggggtgc gaactaaggc c 31

25 <210> 4  
<211> 1732  
<212> DNA  
30 <213> Bos taurus

35 <220>  
<221> misc\_feature  
<222> (1)..(3)  
40 <223> translation start codon

45 <220>  
<221> misc\_feature  
<222> (1468)..(1470)  
50 <223> translation stop codon

55 <220>  
<221> variation  
<222> (689)..(755)  
60 <223> this sequence is deleted in the alternately spliced transcript

65 <220>  
<221> variation

<222> (358)..(358)

<223> cytosine (C) to thymine (T) substitution polymorphism

5

<220>

<221> variation

10

<222> (694)..(695)

<223> adenine (A)-adenine (A) to guanine(G)-cytosine (C) substitution p  
olymorphism  
AA corresponds to the Q allele  
GC corresponds to the q allele

15

20

<220>

<221> primer\_bind

<222> (616)..(635)

25

<223> Primer DgatforAD  
TTCTCCTACCGGGACGTCAA

30

<220>

<221> primer\_bind

35

<222> (743)..(767)

<223> Primer DgatrevAD  
AAGTAGTAGAGATCGCGGTAGGTCA  
reverse primer

40

<220>

45

<221> primer\_bind

<222> (690)..(703)

<223> Primer ForAA (FAM)  
CGTTGGCCTTCTTA  
reverse primer

50

55

<220>

<221> primer\_bind

<222> (688)..(709)

60

<223> Primer DgatADGC (VIC)  
TTGGCCGCCTTACC  
reverse primer

65



<220>

<221> primer\_bind

5 <222> (617)..(636)

<223> Primer DgatforRT66  
TCTCCTACCGGGACGTCAAC

10 <220>

<221> primer\_bind

15 <222> (739)..(760)

<223> Primer DgatrevRT66  
CAGATCGCGGTAGGTCAGGTT  
20 reverse primer

<220>

25 <221> primer\_bind

<222> (676)..(767)

30 <223> Primer DgatforRTless66  
GCTGCTTTGGCAGATCTCTACTACTT  
This primer selectively binds and amplifies a contiguous sequence  
characteristic of the splice variant & generated by the deletion  
bases 689 to 755 of this sequence.

35 <220>

<221> primer\_bind

40 <222> (817)..(833)

<223> Primer DgatrevRTless66  
45 AAGCGCTTTCGGATGCG

<220>

50 <221> primer\_bind

<222> (722)..(734)

55 <223> Primer Dgatwith66 (FAM)  
CCGTGAGCTACCC

60 <220>

<221> primer\_bind

<222> (771)..(785)

65 <223> Primer Dgatless66 (VIC)  
CTTCGCCCCCACCCT

<400> 4	
5	atgggcgacc gcggcggcgc gggcggtccc cggcgccgga ggacggggtc gcggccttcg 60
	atccagggcg gcagtgggcc cggcgagcgc gaagaggagg tgcgggatgt gggcgccgga 120
10	ggggacgcgc cggctccgga cacagacaag gacggagacg tagacgtggg cagcgccac 180
	tgggacctga ggtgtcaccg cctgcaggat tccctgttca gttctgacag tggcttcagc 240
	aactaccgtg gcctcctgaa ttggtgtgtg gtgatgtga tcttaagcaa cgcacggtta 300
15	tttctagaga acctcatcaa gtatggcatc ctggtggacc ccatccaggt ggtgtctctg 360
	ttcctgaagg acccctacag ctggccagct ctgtgcctgg tcattgtggc caatatcttt 420
20	gccgtggctg cgttcacagt ggagaagcgc ctggccgtgg gagctctgac ggagcaggcg 480
	gggctgtctg tgcacggggt caacctggcc accattctct gcttcccagc ggccgtggcc 540
	tttctcctcg agtctatcac tccagtgggc tccgtgtctg ccctgatggg ctacaccatc 600
25	ctcttctcga agctgttctc ctaccgggac gtcaacctct ggtgccgaga gcgcagggct 660
	ggggccaagg ccaaggctgc tttggcaggt aagaaggcca acgggggagc tgcccagcgc 720
30	accgtgagct accccgacaa cctgacctac cgcgatctct actacttctc cttcgccccc 780
	accctgtgct acgagctcaa cttccccgcg tccccccgca tccgaaagcg cttcctgtctg 840
	cggcgactcc tggagatgct gttcctcacc cagctccagg tggggctgat ccagcagtgg 900
35	atggtcccg gcctccagaa ctccatgaag cccttcaagg acatggacta ctcccgcatc 960
	gtggagcgcc tctgaagct ggcggtcccc aaccacctca tctggctcat cttcttctac 1020
40	tggctcttcc actcctgcct gaacgcccgtg gctgagctca tgcagtgttg agaccgcgag 1080
	ttctaccggg actggtggaa ctccgagtc atcacctact tctggcagaa ctggaacatc 1140
	cctgttcaca agtgggtgcat cagacacttc tacaagccca tgctccggcg gggcagcagc 1200
45	aagtgggag ccaggacggc agtgtttctg gcctccgctc tcttccacga gtacctggtg 1260
	agcatcccc tgccgatgtt ccgcctctgg gccttcaccg gcatgatggc gcagatcccc 1320
50	ctggcctgga tagtggggcg cttcttccgc ggcaactacg gcaacgcggc cgtgtggctg 1380
	tcactcatca tcgggcagcc ggtggccgtc ctgatgtacg tccacgacta ctacgtgtctc 1440
	aaccgtgagg cgccgcgagc cggcacctga gcgcctccag gctggccccc tcgtgggtgt 1500
55	tggactgctt tgccgcgctg cctgcggctg gactagagcc tgccccaacc tgggcgcagc 1560
	aggaggaggc ctggctggtg gaagctgcct cctggcctcc accaggcctc tgcctgaagg 1620
60	gcttctcct gccaggggag agcaggcccg acgcagttct ggcccctggg aggtgcccac 1680
	gctctggaaa ccctacagat ctgcaccaag ggtctgaatg tgtcaataaa ga 1732
<210> 5	
65	<211> 489

<212> PRT  
 <213> Bos taurus  
 5  
 <220>  
 <221> VARIANT  
 10  
 <222> (232)..(232)  
 <223> an amino acid substitution (K -> A) caused by a polymorphism at b  
 ases 7224-7225 of the genomic sequence (measured from the adenine  
 15 residue of the translation start codon). Lysine (K) corresponds  
 to the Q allele, alanine (A) corresponds to the q allele.  
 20 <400> 5  
 Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Arg Thr Gly  
 1 5 10 15  
 25 Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu  
 20 25 30  
 30 Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr  
 35 40 45  
 35 Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asp Leu Arg  
 50 55 60  
 40 Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser Asp Ser Gly Phe Ser  
 65 70 75 80  
 Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val Met Leu Ile Leu Ser  
 85 90 95  
 45 Asn Ala Arg Leu Phe Leu Glu Asn Leu Ile Lys Tyr Gly Ile Leu Val  
 100 105 110  
 50 Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro Tyr Ser Trp  
 115 120 125  
 55 Pro Ala Leu Cys Leu Val Ile Val Ala Asn Ile Phe Ala Val Ala Ala  
 130 135 140  
 60 Phe Gln Val Glu Lys Arg Leu Ala Val Gly Ala Leu Thr Glu Gln Ala  
 145 150 155 160  
 Gly Leu Leu Leu His Gly Val Asn Leu Ala Thr Ile Leu Cys Phe Pro  
 165 170 175  
 65 Ala Ala Val Ala Phe Leu Leu Glu Ser Ile Thr Pro Val Gly Ser Val  
 180 185 190

5 Leu Ala Leu Met Val Tyr Thr Ile Leu Phe Leu Lys Leu Phe Ser Tyr  
 195 200 205  
 Arg Asp Val Asn Leu Trp Cys Arg Glu Arg Arg Ala Gly Ala Lys Ala  
 210 215 220  
 10 Lys Ala Ala Leu Ala Gly Lys Lys Ala Asn Gly Gly Ala Ala Gln Arg  
 225 230 235 240  
 15 Thr Val Ser Tyr Pro Asp Asn Leu Thr Tyr Arg Asp Leu Tyr Tyr Phe  
 245 250 255  
 20 Leu Phe Ala Pro Thr Leu Cys Tyr Glu Leu Asn Phe Pro Arg Ser Pro  
 260 265 270  
 25 Arg Ile Arg Lys Arg Phe Leu Leu Arg Arg Leu Leu Glu Met Leu Phe  
 275 280 285  
 Leu Thr Gln Leu Gln Val Gly Leu Ile Gln Gln Trp Met Val Pro Ala  
 290 295 300  
 30 Ile Gln Asn Ser Met Lys Pro Phe Lys Asp Met Asp Tyr Ser Arg Ile  
 305 310 315 320  
 35 Val Glu Arg Leu Leu Lys Leu Ala Val Pro Asn His Leu Ile Trp Leu  
 325 330 335  
 40 Ile Phe Phe Tyr Trp Leu Phe His Ser Cys Leu Asn Ala Val Ala Glu  
 340 345 350  
 45 Leu Met Gln Phe Gly Asp Arg Glu Phe Tyr Arg Asp Trp Trp Asn Ser  
 355 360 365  
 Glu Ser Ile Thr Tyr Phe Trp Gln Asn Trp Asn Ile Pro Val His Lys  
 370 375 380  
 50 Trp Cys Ile Arg His Phe Tyr Lys Pro Met Leu Arg Arg Gly Ser Ser  
 385 390 395 400  
 55 Lys Trp Ala Ala Arg Thr Ala Val Phe Leu Ala Ser Ala Phe Phe His  
 405 410 415  
 60 Glu Tyr Leu Val Ser Ile Pro Leu Arg Met Phe Arg Leu Trp Ala Phe  
 420 425 430  
 65 Thr Gly Met Met Ala Gln Ile Pro Leu Ala Trp Ile Val Gly Arg Phe  
 435 440 445  
 Phe Arg Gly Asn Tyr Gly Asn Ala Ala Val Trp Leu Ser Leu Ile Ile

76

450                      455                      460  
 5    Gly Gln Pro Val Ala Val Leu Met Tyr Val His Asp Tyr Tyr Val Leu  
      465                      470                      475                      480  
 10    Asn Arg Glu Ala Pro Ala Ala Gly Thr  
                                  485  
 15    <210> 6  
      <211> 467  
      <212> PRT  
      <213> Bos taurus  
 20    <400> 6  
 25    Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Arg Thr Gly  
      1                      5                      10                      15  
 30    Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu  
                          20                      25                      30  
 35    Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr  
                          35                      40                      45  
 40    Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asp Leu Arg  
                          50                      55                      60  
 45    Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser Asp Ser Gly Phe Ser  
      65                      70                      75                      80  
 50    Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val Met Leu Ile Leu Ser  
                          85                      90                      95  
      Asn Ala Arg Leu Phe Leu Glu Asn Leu Ile Lys Tyr Gly Ile Leu Val  
                          100                      105                      110  
 55    Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro Tyr Ser Trp  
                          115                      120                      125  
      Pro Ala Leu Cys Leu Val Ile Val Ala Asn Ile Phe Ala Val Ala Ala  
                          130                      135                      140  
 60    Phe Gln Val Glu Lys Arg Leu Ala Val Gly Ala Leu Thr Glu Gln Ala  
      145                      150                      155                      160  
 65    Gly Leu Leu Leu His Gly Val Asn Leu Ala Thr Ile Leu Cys Phe Pro  
                          165                      170                      175

77

	Ala	Ala	Val	Ala	Phe	Leu	Leu	Glu	Ser	Ile	Thr	Pro	Val	Gly	Ser	Val	
				180						185					190		
5	Leu	Ala	Leu	Met	Val	Tyr	Thr	Ile	Leu	Phe	Leu	Lys	Leu	Phe	Ser	Tyr	
			195					200					205				
10	Arg	Asp	Val	Asn	Leu	Trp	Cys	Arg	Glu	Arg	Arg	Ala	Gly	Ala	Lys	Ala	
		210					215					220					
15	Lys	Ala	Ala	Leu	Ala	Asp	Leu	Tyr	Tyr	Phe	Leu	Phe	Ala	Pro	Thr	Leu	
	225				230						235					240	
	Cys	Tyr	Glu	Leu	Asn	Phe	Pro	Arg	Ser	Pro	Arg	Ile	Arg	Lys	Arg	Phe	
					245					250					255		
20	Leu	Leu	Arg	Arg	Leu	Leu	Glu	Met	Leu	Phe	Leu	Thr	Gln	Leu	Gln	Val	
				260					265						270		
25	Gly	Leu	Ile	Gln	Gln	Trp	Met	Val	Pro	Ala	Ile	Gln	Asn	Ser	Met	Lys	
		275					280						285				
30	Pro	Phe	Lys	Asp	Met	Asp	Tyr	Ser	Arg	Ile	Val	Glu	Arg	Leu	Leu	Lys	
		290					295					300					
35	Leu	Ala	Val	Pro	Asn	His	Leu	Ile	Trp	Leu	Ile	Phe	Phe	Tyr	Trp	Leu	
	305				310					315						320	
	Phe	His	Ser	Cys	Leu	Asn	Ala	Val	Ala	Glu	Leu	Met	Gln	Phe	Gly	Asp	
					325					330					335		
40	Arg	Glu	Phe	Tyr	Arg	Asp	Trp	Trp	Asn	Ser	Glu	Ser	Ile	Thr	Tyr	Phe	
				340					345						350		
45	Trp	Gln	Asn	Trp	Asn	Ile	Pro	Val	His	Lys	Trp	Cys	Ile	Arg	His	Phe	
		355					360						365				
50	Tyr	Lys	Pro	Met	Leu	Arg	Arg	Gly	Ser	Ser	Lys	Trp	Ala	Ala	Arg	Thr	
		370					375					380					
55	Ala	Val	Phe	Leu	Ala	Ser	Ala	Phe	Phe	His	Glu	Tyr	Leu	Val	Ser	Ile	
	385				390						395					400	
	Pro	Leu	Arg	Met	Phe	Arg	Leu	Trp	Ala	Phe	Thr	Gly	Met	Met	Ala	Gln	
					405					410					415		
60	Ile	Pro	Leu	Ala	Trp	Ile	Val	Gly	Arg	Phe	Phe	Arg	Gly	Asn	Tyr	Gly	
				420					425					430			
65	Asn	Ala	Ala	Val	Trp	Leu	Ser	Leu	Ile	Ile	Gly	Gln	Pro	Val	Ala	Val	
		435						440					445				

Leu Met Tyr Val His Asp Tyr Tyr Val Leu Asn Arg Glu Ala Pro Ala  
450 455 460

5 Ala Gly Thr  
465

10 <210> 7  
<211> 12  
<212> DNA  
15 <213> Bos taurus

20 <220>  
<221> variation  
<222> (7)..(7)  
25 <223> thymidine (T) to guanine (G) substitution polymorphism

30 <400> 7  
cagtgctagg gg 12

35 <210> 8  
<211> 10  
<212> DNA  
40 <213> Bos taurus

45 <220>  
<221> variation  
<222> (6)..(6)  
50 <223> thymidine (T) to guanine (G) substitution polymorphism

55 <400> 8  
gcattgcgct 10

60 <210> 9  
<211> 11  
<212> DNA  
65 <213> Bos taurus

<220>  
<221> variation  
5 <222> (6)..(6)  
<223> thymidine (T) to cytosine (G) substitution polymorphism  
10  
<400> 9  
taccctggga c 11  
15 <210> 10  
<211> 11  
<212> DNA  
20 <213> Bos taurus  
25 <220>  
<221> variation  
<222> (6)..(6)  
30 <223> adenine (A) to guanine (G) substitution polymorphism  
35 <220>  
<221> variation  
<222> (6)..(6)  
40 <223> adenine (A) to guanine (G) substitution polymorphism  
45 <400> 10  
ctcttagcag c 11  
50 <210> 11  
<211> 10  
<212> DNA  
55 <213> Bos taurus  
60 <220>  
<221> variation  
<222> (5)..(5)  
65 <223> guanine (G) to adenine (A) substitution polymorphism



	<400> 11	
5	acaggcaact	10
	<210> 12	
	<211> 11	
10	<212> DNA	
	<213> Bos taurus	
15	<220>	
	<221> variation	
20	<222> (6)..(6)	
	<223> cytosine (C) to thymine (T) substitution polymorphism	
25	<400> 12	
	tgtctctgtt c	11
30	<210> 13	
	<211> 14	
35	<212> DNA	
	<213> Bos taurus	
40	<220>	
	<221> variation	
45	<222> (7)..(8)	
	<223> adenine (A)-adenine (A) to guanine(G)-cytosine (C) substitution p olymorphism	
50	AA corresponds to the Q allele	
	GC corresponds to the q allele	
55	<400> 13	
	ggtaagaagg ccaa	14
	<210> 14	
60	<211> 11	
	<212> DNA	
65	<213> Bos taurus	

<220>  
<221> variation  
5 <222> (6)..(6)  
<223> guanine (G) to adenine (A) substitution polymorphism  
10  
<400> 14  
gcggtgagga t 11  
15 <210> 15  
<211> 16  
<212> DNA  
20 <213> Bos taurus  
25 <220>  
<221> variation  
<222> (6)..(7)  
30 <223> guanine (G)-guanine (G) to adenine (A)-cytosine (C) substitution  
polymorphism  
GG-AC  
35  
<400> 15  
gggggggggg gactct 16  
40  
<210> 16  
<211> 11  
45 <212> DNA  
<213> Bos taurus  
50  
<220>  
<221> variation  
55 <222> (6)..(6)  
<223> adenine (A) to guanine (G) substitution polymorphism  
60  
<400> 16  
gagtgcctg c 11  
65 <210> 17

<211> 11  
<212> DNA  
5 <213> Bos taurus

10 <220>  
<221> variation  
<222> (6)..(6)  
15 <223> cytosine (C) to thymidine (T) substitution polymorphism

20 <400> 17  
ggacgcgtgg g 11

<210> 18  
25 <211> 11  
<212> DNA  
<213> Bos taurus  
30

<220>  
35 <221> variation  
<222> (6)..(6)  
<223> guanine (G) to adenine (A) substitution polymorphism  
40

<400> 18  
45 ggtgggggtg g 11

<210> 19  
50 <211> 11  
<212> DNA  
<213> Bos taurus

55 <220>  
<221> variation  
60 <222> (6)..(6)  
<223> ctosine (C) to thymidine (T) substitution polymorphism  
65

<400> 19

83

ctgggcgcag c 11

5 <210> 20  
<211> 14  
<212> DNA

10 <213> Bos taurus

15 <400> 20  
cgttggcctt ctta 14

<210> 21

20 <211> 14  
<212> DNA

25 <213> Bos taurus

<400> 21  
30 ttggccgcct tacc 14

35

**CLAIMS:**

1. A method of determining genetic merit of a bovine with respect to milk composition and volume which comprises the step of determining the DGAT1 genotypic state of said bovine.  
5
2. A method as claimed in claim 1, wherein the genotypic state is determined with respect to DNA, mRNA and/or protein obtained from said bovine by direct or indirect methods.  
10
3. A method as claimed in claim 1, wherein the genotypic state is determined by the presence of at least one nucleotide difference from the nucleotide sequence encoding bovine DGAT1 (SEQ ID NO: 1) either by direct or indirect methods.
- 15 4. A method as claimed in claim 3, wherein the genotypic state is determined by the presence of one or more polymorphisms of SEQ ID NOs: 7 to 19 either by direct or indirect methods.
- 20 5. A method as claimed in claim 4, wherein the genotypic state is determined by detecting the presence of the K232A polymorphism (SEQ ID NO: 13), either by direct or indirect methods.
- 25 6. A method of selecting a bovine having a desired DGAT1 genotypic state comprising determining the genotypic state according to any one of claims 1 to 5 and selecting said bovine on the basis of said determination.
7. A bovine selected by the method of claim 6.
8. A method of identifying a bovine which possesses a genotype indicative of altered milk production traits, said method comprising:  
30 obtaining a nucleic acid sample from said bovine and identifying a polymorphism selected from the group comprising SEQ ID NOs: 7 to 19 of the bovine DGAT1 gene,  
wherein the presence of said polymorphism is associated with altered milk  
35 production traits.

9. A method as claimed in claim 8, wherein said altered milk production traits comprise an increase in milk volume and/or an increased protein to fat ratio of the milk composition or a decrease in milk volume and/or an increase in fat content of the milk composition.

5

10. A method as claimed in claim 8, wherein the polymorphism is K232A (SEQ ID NO: 13), wherein the presence or absence of an alanine is associated with the altered milk production traits as claimed in claim 9.

10

11. A method as claimed in claim 8, further comprising the step of amplifying said bovine DGAT1 gene sequence.

12. A method as claimed in claim 11, wherein primers selected from the group consisting of (SEQ ID NOs: 20 and 21) are used in said amplification.

15

13. A primer suitable for use in detecting a polymorphism selected from the group comprising SEQ ID NOs: 7 to 19 of the bovine DGAT1 gene, said primer consisting of a nucleotide sequence having about at least 12 contiguous bases of SEQ ID NOs. 1, 3 or 4.

20

14. A bovine identified by the method of any one of claims 8 to 12.

15. A bovine as claimed in claim 7 or 14, comprising a bull.

25

16. Semen produced by a bovine as claimed in claim 15.

17. A bovine as claimed in claim 7 or 14, comprising a cow.

18. Milk produced by a bovine as claimed in claim 17.

30

19. Milk as claimed in claim 18, comprising one or more altered characteristics selected from the group consisting of increased % protein; increased total protein content; increased protein to fat ratio; decreased total fat content; increased fat softness; increase unsaturated fat content; increased or decreased solid fat content; and increased or decreased total volume; when compared to milk produced by a non-genotyped bovine.

35

20. Milk as claimed in claim 18, which has a decreased solid fat content, and increased fat softness and is advantageous for use in dairy product processing plants.
- 5 21. A dairy product made from the milk as claimed in any one of claims 18 to 20.
22. A dairy product made from the milk as claimed in claim 20, wherein said product provides a beneficial health effect.
- 10 23. The use of the DGAT1 gene sequence (SEQ ID NOs: 1, 3 and 4) or a fragment or variant thereof, in the identification of one or more molecular DNA markers useful in genotyping and/or selecting a bovine according to the methods of any one of claims 1 to 6 and 8 to 12.
- 15 24. The use of one or more polymorphism sequences selected from the group consisting of SEQ ID NOs: 7 to 19 in a method of identification and selection of a bovine having at least one of said polymorphisms in its DGAT1 gene.
- 20 25. The use of a probe in the methods of genotyping according to any one of claims 1 to 6 and 8 to 12, wherein the probe is selected from any 5 or more contiguous nucleotides of the DGAT1 sequence of SEQ ID NOs: 1, 3 or 4, which is sufficiently complementary with said nucleic acid sequence so as to bind thereto under stringent conditions.
- 25 26. A kit for genotyping a bovine with respect to milk composition and volume associated with DGAT1, comprising a primer or probe as claimed in claim 13 or 25.
27. An isolated nucleic acid molecule having the sequence of SEQ ID NOs: 1 or 4 with or without one or more polymorphisms selected from the group consisting of SEQ  
30 ID Nos: 7 to 19, or a fragment or variant thereof.
28. An isolated nucleic acid molecule comprising a DNA molecule having in whole or in part the nucleotide sequence of SEQ ID NOs: 1, 3 or 4 or which varies from the sequence due to the degeneracy of the genetic code, or a nucleic acid strand capable of  
35 hybridising with said nucleic acid molecule under stringent hybridisation conditions.

29. An isolated mRNA transcribed from DNA having a sequence which corresponds to a nucleic acid molecule as claimed in claim 27 or 28.
30. A recombinant cloning vector comprising the DNA molecule of claim 27 or 28.
- 5 31. A prokaryotic or eukaryotic cell containing the cloning vector of claim 30.
32. A transfected cell line which expresses a protein encoded by the nucleic acid molecule of claim 27 or 28.
- 10 33. A diagnostic kit useful in detecting a DNA molecule as claimed in claim 27 or 28, comprising a first and second primer for amplifying the DNA, the primers being complementary to nucleotide sequences of the DNA upstream and downstream, respectively, of a polymorphism in the portion of the DNA encoding DGAT1 which results in altered relative milk lipid and protein production and milk volume, wherein at least one of the nucleotide sequences is selected to be from a non-coding region of the DGAT1 gene.
- 15 34. A kit as claimed in claim 33, further comprising a third primer complementary to a polymorphism selected from the group consisting of of SEQ ID NO: 7 to 19 located on the DGAT1 gene.
- 20 35. A process for producing a protein encoded by a DNA molecule of claim 27 or 28, comprising the steps:
- 25 a) preparing a DNA fragment including a nucleotide sequence which encodes the protein;
- b) incorporating the DNA fragment into an expression vector to obtain a recombinant DNA molecule which includes the DNA fragment and is capable of undergoing replication;
- 30 c) transforming a host cell with the recombinant DNA molecule to produce a transformant which can express the protein;
- d) culturing the transformant to produce the protein; and
- e) recovering the protein from resulting cultured mixture.
- 35 36. A purified protein having an amino acid sequence of SEQ ID NOs: 2, 5 or 6 or a fragment or variant thereof having biological activity of DGAT1.



37. An antibody raised against the protein as claimed in claim 36.
38. A method for the modulation of milk production and content in a bovine by modulating the activity of the DGAT1 protein.
- 5 39. A method as claimed in claim 38, comprising administering to the bovine an effective amount of a nucleic acid molecule substantially complementary to at least a portion of mRNA encoding the bovine DGAT1 protein and being of sufficient length to sufficiently reduce expression of said DGAT1.
- 10 40. An isolated antisense nucleic acid molecule for use in a method as claimed in claim 39.
41. A method of modulating milk production and/or milk content in a lactating  
15 bovine, including administering to the bovine an effective amount of a nucleic acid molecule having ribozyme activity and a nucleotide sequence substantially complementary to at least a portion of mRNA encoding a bovine DGAT1 and being of sufficient length to bind selectively thereto to sufficiently reduce expression of said DGAT1.
- 20 42. A method of inhibiting the activity of bovine DGAT1 in a lactating bovine so as to modulate milk production and/or milk solids content, comprising administering an effective amount of an antibody to the relevant DGAT1.
- 25 43. A method of modulating milk production and/or milk solids content by raising an autoantibody to a bovine DGAT1 in the bovine.
44. A transgenic animal having cells that contain a transgene, which transgene is introduced into the animal or an ancestor of the animal at a prenatal stage, wherein  
30 said transgene comprises a bovine cDNA comprising the nucleotide sequence of SEQ ID NO: 4 or an appropriate variant or subsequence thereof.
45. A transgenic non-human animal wherein said endogenous DGAT1 gene has been knocked out and replaced with a transgene.
- 35 46. A transgenic non-human animal, wherein additional copies of the gene encoding DGAT1 are inserted as a transgene.

47. A transgenic non-human animal as claimed in claim 45 comprising a transgenic bovine having a genome lacking a gene encoding a protein having biological activity of DGAT1.

5


48. A transgenic non-human animal comprising a mouse having a genome containing a gene encoding a bovine protein having biological activity of DGAT1.

49. A transgenic bovine having a gene encoding a bovine protein having biological activity of bovine DGAT1 and heterologous nucleotide sequence antisense to the gene.

10

50. A transgenic bovine having a genome which includes additional copies of a gene encoding a protein having biological activity of DGAT1 or copies of a gene encoding a protein having biological activity of DGAT1 under control of a high expression promoter.

15



### FIGURE 1

2/34

<222> (29)..(31)

<223> bases 1 to 3 of the Kozak recognition sequence. See the genomic sequence from the start codon for bases 4 to 7 of the Kozak recognition sequence or the DGAT1 cDNA for the complete recognition sequence.

acttggccgc ggcggggtgc gaactaaggc c

**FIGURE 2a**

3/34

<210> 1

<211> 11771

<212> DNA

<213> Bos taurus

<220>

<221> CDS

<222> (1)..(191)

<223> Exon 1 CDS, determined by alignment with an amino acid sequence deduced from the cDNA

<220>

<221> misc\_signal

<222> (1)..(4)

<223> these bases correspond to bases 4 to 7 of the Kozak recognition sequence. See DGAT1 cDNA for the complete recognition sequence.

<220>

<221> CDS

<222> (3809)..(3896)

<223> Exon 2, determined by alignment with an amino acid sequence deduced from the cDNA

<220>

<221> CDS

<222> (5840)..(5880)

<223> Exon 3, determined by alignment with an amino acid sequence deduced from the cDNA

FIGURE 2b

4/34

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (5960)..(6045)

&lt;223&gt; Exon 4, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (6138)..(6190)

&lt;223&gt; Exon 5, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (6406)..(6511)

&lt;223&gt; Exon 6, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (6601)..(6714)

&lt;223&gt; Exon 7, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (6815)..(6889)

&lt;223&gt; Exon 8, determined by alignment with an amino acid sequence deduced from the cDNA

FIGURE 2b continued

5/34

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (6824)..(6889)

&lt;223&gt; this sequence is deleted from the alternately spliced transcript.

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (6960)..(7063)

&lt;223&gt; Exon 9, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (7154)..(7192)

&lt;223&gt; Exon 10, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (7271)..(7312)

&lt;223&gt; Exon 11, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (7386)..(7430)

&lt;223&gt; Exon 12, determined by alignment with an amino acid sequence deduced from the cDNA

FIGURE 2b continued

6/34

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (7505)..(7617)

&lt;223&gt; Exon 13, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (7705)..(7770)

&lt;223&gt; Exon 14, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (7858)..(7945)

&lt;223&gt; Exon 15, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (8027)..(8089)

&lt;223&gt; Exon 16, determined by alignment with an amino acid sequence deduced from the cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (8162)..(8314)

&lt;223&gt; Exon 17 CDS, determined by alignment with an amino acid sequence deduced from the cDNA

**FIGURE 2b continued**



7/34

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (8315)..(8317)

&lt;223&gt; translation stop codon

&lt;220&gt;

&lt;221&gt; polyA\_site

&lt;222&gt; (8572)..(8578)

&lt;223&gt;

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (626)..(626)

&lt;223&gt; thymidine (T) to guanine (G) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (3512)..(3512)

&lt;223&gt; thymidine (T) to guanine (G) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (4040)..(4040)

&lt;223&gt; thymidine (T) to cytosine (G) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (4963)..(4963)

&lt;223&gt; adenine (A) to guanine (G) substitution polymorphism

**FIGURE 2b continued**

8/34

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (5003)..(5003)

&lt;223&gt; guanine (G) to adenine (A) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (5997)..(5997)

&lt;223&gt; cytosine (C) to thymine (T) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (6829)..(6830)

<223> adenine (A)-adenine (A) to guanine(G)-cytosine (C) substitution polymorphism  
AA corresponds to the Q allele  
GC corresponds to the q allele

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (6892)..(6892)

&lt;223&gt; guanine (G) to adenine (A) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (7224)..(7225)

<223> guanine (G)-guanine (G) to adenine (A)-cytosine (C) substitution polymorphism  
GG-AC

FIGURE 2b continued

9/34

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (7438)..(7438)

&lt;223&gt; adenine (A) to guanine (G) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (7456)..(7456)

&lt;223&gt; cytosine (C) to thymidine (T) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (7987)..(7987)

&lt;223&gt; guanine (G) to adenine (A) substitution polymorphism

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (8402)..(8402)

&lt;223&gt; ctosine (C) to thymidine (T) substitution polymorphism

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (9434)..(9434)

&lt;223&gt; ambiguous nucleotide

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (9496)..(9496)

&lt;223&gt; ambiguous nucleotide

**FIGURE 2b continued**

10/34

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (10402)..(10417)

&lt;223&gt; ambiguous nucleotides

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6579)..(6601)

<223> Primer 17F  
CCTGAGCTTGCCTCTCCCACAGT

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (7036)..(7058)

<223> Primer 18R  
CCAGGAGTCGCCGCAGCAGGAAG  
reverse primer

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (7280)..(7303)

<223> Primer 6F  
CCGGCCATCCAGAACTCCATGAAG

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (7585)..(7605)

<223> Primer AW446985 dn1  
TAGAACTCGCGGTCTCCAAAC  
reverse primer

FIGURE 2b continued

11/34

<220>  
<221> primer\_bind  
<222> (8222)..(8244)  
<223> Primer InsUp1  
TGGCTGTCACTCATCATCGGGCA

<220>  
<221> primer\_bind  
<222> (8566)..(8589)  
<223> Primer 14R2  
TTGCACAGCACTTTATTGACACA

<220>  
<221> primer\_bind  
<222> (6813)..(6830)  
<223> Primer SNP1\_FAM  
AGC TTT GGC AGG TAA GGC

<220>  
<221> primer\_bind  
<222> (6813)..(6830)  
<223> Primer SNP1\_HEX  
AGC TTT GGC AGG TAA GAA

<220>  
<221> primer\_bind  
<222> (6831)..(6844)  
<223> Primer SNP1\_2P  
GGC CAA CGG GGG AG

FIGURE 2b continued

12/34

<220>  
<221> primer\_bind  
<222> (7424)..(7438)  
<223> Primer SNP2\_FAM  
GCT GGC GGT GAG TGA

<220>  
<221> primer\_bind  
<222> (7424)..(7438)  
<223> Primer SNP2\_HEX  
GCT GGC GGT GAG TGG

<220>  
<221> primer\_bind  
<222> (7439)..(7453)  
<223> Primer SNP2\_2P  
CCT GCT GGG TGG GGA

<220>  
<221> primer\_bind  
<222> (7442)..(7456)  
<223> Primer SNP3\_FAM  
GCT GGG TGG GGA CGC

<220>  
<221> primer\_bind  
<222> (7442)..(7456)  
<223> Primer SNP3\_HEX  
GCT GGG TGG GGA CGT

FIGURE 2b continued

13/34

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (7457)..(7470)

<223> Primer SNP3\_P  
GTG GGG GCG GGT GG

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (8388)..(8402)

<223> Primer SNP4\_FAM  
TGC CCC AAC CTG GGT

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (8388)..(8402)

<223> Primer SNP4\_HEX  
TGC CCC AAC CTG GGC

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (8403)..(8417)

<223> Primer SNP4\_2P  
GCA GCA GGA GGA GGC

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6811)..(6830)

<223> Primer Dgat 21  
GTAGCTTTGGCAGGTAAGAA

FIGURE 2b continued

14/34

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6965)..(6984)

<223> Primer Dgat 22  
GGGGCGAAGAGGAAGTAGTA  
reverse primer

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6613)..(6632)

<223> Primer Dgat 23  
TGGCCCTGATGGTCTACACC

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6829)..(6850)

<223> Primer Dgat 24B  
GGGCAGCTCCCCCGTTGGCCGC  
reverse primer

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6651)..(6670)

<223> Primer DgatforAD  
TTCTCCTACCGGACGTCAA

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6871)..(6890)

<223> Primer ReverseNZ  
CCGCGGTAGGTCAGGTTGTC  
reverse primer

FIGURE 2b continued



15/34

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6825)..(6838)

<223> Probe ForAA (FAM)  
CGTTGGCCTTCTTA

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6823)..(6836)

<223> Probe DgatADGC (VIC)  
TTGCCGCCTTACC

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6651)..(6670)

<223> Primer DgatforAD  
TTCTCCTACCGGGACGTCAA

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6878)..(6972)

<223> Primer DgatrevAD  
AAGTAGTAGAGATCGCGGTAGGTCA  
reverse primer

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6825)..(6838)

<223> Probe ForAA (FAM)  
CGTTGGCCTTCTTA

FIGURE 2b continued

16/34

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6823)..(6836)

<223> Probe DgatADGC (VIC)  
TTGGCCGCCTTACC

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6652)..(6671)

<223> Primer DgatforRT66  
TCTCCTACCGGACGTCAAC

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6874)..(6964)

<223> Primer DgatrevRT66  
GAGATCGCGGTAGGTCAGGTT  
reverse primer

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6711)..(6972)

<223> Primer DgatforRTless66, GCTGCTTTGGCAGATCTCTACTACTT  
This primer was designed to selectively bind and amplify the cDNA  
splice variant. The corresponding binding site in this genomic s  
equence comprises bases 6711 to 6715, 6815 to 6823 and 6960 to 69  
72.

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (7022)..(7038)

<223> Primer DgatrevRTless66  
AAGCGCTTTCGGATGCG  
reverse primer

FIGURE 2b continued

17/34

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6857)..(6870)

<223> Probe Dgatwith66 (FAM)  
CCGTGAGCTACCC

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (6976)..(6990)

<223> Probe Dgatless66 (VIC)  
CTTCGCCCCACCCCT

&lt;400&gt; 1

atg ggc gac cgc ggc ggc gcg ggc ggc tcc cgg cgc cgg agg acg ggg	48
Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Arg Thr Gly	
1 5 10 15	
tcg cgg cct tcg atc cag ggc ggc agt ggg ccc gcg gca gcg gaa gag	96
Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu	
20 25 30	
gag gtg cgg gat gtg ggc gcc gga ggg gac gcg ccg gtc cgg gac aca	144
Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr	
35 40 45	
gac aag gac gga gac gta gac gtg ggc agc ggc cac tgg gac ctg ag	191
Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asp Leu Arg	
50 55 60	
gtagcgggtgc gcgtgacccc taacctttga cccctgatac gggggccctg cgacccaacc	251
tggtggccca ggcctgtcgg cggcagctcg ggctcgagtc cgagagtctg gcgcctggac	311
cttggtgcac agctgtgccc ctcgggcctc cacggggaaa cttagcggga gggtggggc	371
ggaggggtctc ctgcccggaa caccaggtta cgggggccga ggggagggca gcggtcaac	431
ttctagacgc cctccctctg ccttcttttg gtgggttctg aagctttccc agggtagacc	491
cactacgcac agtgtctct acctggaagg agatacaggg gtccttctctg agggctatga	551
gggggtgcctt gtgggttgat aaagctcccg ggggaggagg gtggaccggc ggagaacaga	611
ggcaggggca gtgctagggg atttctcatc cctcgagac cctccagaga atggtcttca	671
caaaggtccc tcacccgtca cccggcgatt gactggccta ggatcctgct tattaccage	731
acaaatggct gctctagggc caaagtgggt cctgtaatgg gaccctcacc cctggttggg	791

FIGURE 2b continued

## 18/34

gtacagggga ggagttggaa gtgcgcacac ccacaggtgg gcgccctgct tagctgaagg	851
actgatggga aggagttggg ggagcaagct gcggctgaaa gggaggatct gacccacgtg	911
ggcatcagct aagtcctgct ggctgcctcc aggcgttccc ttggccatcc tccacgcccc	971
tccccgggg cctgaccttc atcctggtca agggctctca ggggctctgg ttttgggac	1031
agctccagag ctagaggtta tcaaggagga agtgggcaac aggtcagtca gcaaggattt	1091
gctatcttca ctgggtgctg tggggagggg agggacaagg gcagttgggg tgcaggcact	1151
gtccctgccc ttggggggca cacagttcac ctgagagata agatagccgc agccctgaag	1211
agtgagagca aaggtcaggc acagagttca ggatgacacc aggggagggg ggctctgtga	1271
ggggcactgg cttcctacag gccccagggt gtccctgaggg ggccgctgca aaggccagga	1331
ggccacaggg cccctctgcc cactcctggg gaactggatt tggggtcact ttgtatgagg	1391
tgggggcggg taccagcttt gggccaagct gtcacctgg atgggccatc acttgccctgc	1451
tctgtatagg ccagatggcc agaagctgct cctgtcctgt tgatggccca tcctcgaggt	1511
ctggacctc gggaagagga gcagttgggt gcagggatgg gccaccggag accctcctga	1571
ctccaggac acgcagctgt gtgtgcctgt cccaagcca catgccacat ggctaggggc	1631
ctcctggggc agggctgggc attggtctgg ctactcttgg tatcgctat ggcttccctg	1691
ctccacgtc atcatcctcc cacctctgcc tcctgcctg ttctctcttt tctctcagg	1751
cccttccgga catttccctgc tcacctaggt ctgggcaggc ggggtcaggt gccgggtgtg	1811
agctcactcc ttccggcagc aaggtgtagc tatgtgccg aaggaaggcc gctgctgttg	1871
cctgcctct gagtgcatcc cttccaggtc ctccacactc ccctgtgcc cgacacctgg	1931
tgcgtccttc agccattggt tcatgtgtcc tcaggcaca gctttctagt ccagagcctc	1991
taggctgggt gcaggaagt ctgaggaagt ggcagccggg aggcgagctg gcacctgtc	2051
cctccttgtt ctgtccgtcc ctgccccctg accgtatggc cccgcatgtg tgatccccac	2111
ttggggctgt gcctctgggc aagttgggaa gcttgggtgag cctcattttc atgtgccgc	2171
ctcccagtac tgatgtgcag gttgaatgag gtgccaaactg taatgagttg gaatggccct	2231
gctggctggg tgggactggg gagcaggtgg gggccgctgg ggggcacaga ggacacccca	2291
gtgcctcagt caggagaggg gtgacagaga agctctgggt gaggcccccac ctccactctg	2351
gccatggctg ctgccctttg gtccactgca gtgaactgtg ccatggggct ggacctctgt	2411
ggggattggt gggcagtggt ctttcttccc gcttggggcc tctgacctct gggggcaggg	2471
cgctgcccgg gtgggacagt cggaaggctg gttaggggac ctgaggggtc tgtgtggtgg	2531
ctgggggcag gcctcaggaa tttgacagca gggatctgga aaagctttaa taacattatt	2591

FIGURE 2b continued

## 19/34

tgttgctcagg attgggaaat gctccccctcc cccctcccc tctttcatct tagagactgc	2651
tgcacatctg gtcagtgtgg tcttcttggg ggcccccaag gtggcagggg tcacactgtt	2711
atgaaaccgt cccctgggta tgtggtgcag acatgcacat gcagatgggtg attggcaggt	2771
tgtagcatga ggtggctttg ggacggttcc agtgacagtg agtgggctgg atctgggggg	2831
ttctgggcag gtccatcaag cggatacccc cacagactgt cctcttgga tagttgggcc	2891
tgggagccct gcttgcttg ccaaaaggca ggcgcagagt catgaagaag agggcttggg	2951
ggctcagagc cccactgtgt gtgcagccca ggggtggacct ggaggagggtg cgtgggcagg	3011
ctgggccggc cgggcgggg ggtggggggg cctggtgtga aagggaacca gggccagact	3071
gtcagcgtg cctggctgag gatgctggca cctgtcctc cccagccgtc tgtctcctgg	3131
gtgcagccat ctgagtgtg accccagccg cccctggagg ctggctgttc tcctgtgccc	3191
tattgctggg gacatgtgtc cacaggaggg aaagggaagc cccggcctct ccccttaca	3251
aactggaggc cttgtcfaat gccctggatg gctcctgtg ggcagggttg ttggtgggag	3311
gtggggctgc tgettagaac ccgccagcgg gcttgggctt gggctgagct gcacccctcc	3371
acctctgcct ccagctgagg gttggttcc atctccacca ggcccagcac tgggcacagg	3431
gctctcagag gcaggctctg aaagtccct gctggcttct gcagtggact ccaggcgccg	3491
agccccagg gggctcgc atgcgtcacc ctgcgaagcc acgtgaaggc tgggtcctcc	3551
cctccggaag ggccaaatgc agggcatggg tggtttgaat ggtggcccct gggctccccg	3611
gagggaccag ctgctgtgag ggccgcccc tcccacttc cgtcttgcat caccagctcc	3671
tgtggcactc cccacgcccc gtccccagt gggagcggca ggccccgggt ggctctgccc	3731
gcggaggggg atgtgtgggc ggcggggtg ccttgcctgc agatgctctg ccccgagtgt	3791
ccgtctccgc tctccag g tgt cac cgc ctg cag gat tcc ctg ttc agt tct	3842
Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser	
65 70 75	
gac agt ggc ttc agc aac tac cgt ggc atc ctg aat tgg tgt gtg gtg	3890
Asp Ser Gly Phe Ser Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val	
80 85 90	
atg ctg gtacgtagag tgacacctg gagcaagggt cctgacggcc ggggggccat	3946
Met Leu	
gggtcttct ccagggtag gtgtctgtac ttgtgtagct gtggtgaatg gagctctgtg	4006
ctggcggtg gggtccttg agcagccgta cctggggacc ctaccgggag catgtctatg	4066
ccgtccctgc tgaatccag gagatgcctg cagagggcag cctgggagcc tctgagctgg	4126
ggtctgcgcc ccagggggca ctggagtctc cccagggggc gagagagagt aggcagggat	4186

FIGURE 2b continued

## 20/34

```

ggctctgggtgg ccctgggtgg gggatggctg ctccgtgggc ccaggccctc cctggcagca 4246
caggtgagtg gtcttggggg tccacgtaga acttcctctt ctgttccaaa ttgccctcat 4306
gggtgcggca tgcttgggtg aacctggggg agcaggggtga ggacatgctt ctcagcccag 4366
cccacagctc caggccacac tctgcaggac tctggccctt ccctcagccc tggagggagc 4426
aggactggag tcctgtgtcc gccttgctct gacctggccg aggccactgc tgtggggccc 4486
cagcaggcct gccacgaga aggtggagtg cagggaaccc aggggcagcc ttcaggggtg 4546
ggcaggggtga ggcccactg gggccagccc caccgctcag tgctgatgtg gcgcgaggcc 4606
ttcgcctctc cagctgacgt gtctgcctgc cctgggtgtg gctccagagg ctgcctgtgt 4666
accagggggc cccacgcttc tgtttgtgtt tctgggcagt cccctgggga gcggtggggg 4726
ctgtgtgccca gtccagaccc agtagtccac gcgtccttgt ctctggaggc cgtggcttgt 4786
ccaggactgt ggcaaggtgg tcgtgcaggg caggccctca gcagcctgtc tgttctcctg 4846
cagccccag cctcctggcc ctttggtgca cccacaaagc tccccctcc cccaggagct 4906
ggggccgcct gctgcctcct ctccgcagcc tgggcttcca ggtggctggg cctcttagca 4966
gtccaactc ttgcctgtgg tgggctctca ggacaggcaa ctgccagtcg gcagacattg 5026
caggaccag tggtctctgg taagctggct ggttaggtgt ttagctgggg gatggtgtgg 5086
caggtggccc ctgcctctct gagcctgtca cctcctcggg aagccttctg ggtgggggac 5146
tccaccctatg tcgcctggag aagcatcact tttccacaga gccttctgca acccccgtgg 5206
ggcctgagcc tgggggtggg gaggtgggtg cccctgtctc tgacagagcc agccaggcat 5266
ctggccccag gccactggca agagctcgtt gtgttggggg atctgtcctt tgctgtctgt 5326
gcaggagcgg ccgaggcagg cgggggcgtg agtaggggtg gagaccagc cccagcttcc 5386
ccagccctc aggaccggcc tgctctttcc caccaccca ccaagtgcgt gggcacaccc 5446
cgctgtgag gatggggccg gttggcaggg cggagccctg ggaggggtggc agtgccggcg 5506
gcaggcttg acttactgg ggcttgggtt gtctcgtgtg gccagggcg ctgaccgct 5566
tggtgggacg gacggccgt gggcagcagg tttcttctgc cacggtggca caggcacctg 5626
gggttgtgtt tgctccagg cgggcggggg ctgcgtgcc ctgcgcaggc acataggccg 5686
tgggtgggga gtctcagagc ttggcgtgag gtcccacagg gctgggcctg caggatggag 5746
gccactgtcc tgagctgcag gtgctggcag gagctggggt gggcgttctg gggccgtggc 5806
tgacagcgtt atgtccctct ctctctatcg cag atc tta agc aac gca cgg tta 5860
                                Ile Leu Ser Asn Ala Arg Leu
                                95                                100

ttt cta gag aac ctc atc aa gtgagtgggc ccgggcctgc cccagccct 5910
Phe Leu Glu Asn Leu Ile Lys
                                105

```

FIGURE 2b continued

## 21/34

gccacctcac cctcgccta cacagacct caccacctg cgtctgcag g tat ggc 5966  
Tyr Gly

atc ctg gtg gac ccc atc cag gtg gtg tct ctg ttc ctg aag gac ccc 6014  
Ile Leu Val Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro  
110 115 120 125

tac agc tgg cca gct ctg tgc ctg gtc att g gtgagctggg tgcccaggag 6065  
Tyr Ser Trp Pro Ala Leu Cys Leu Val Ile  
130 135

gcctcaggcc ggcggtgggt gggacagggc tgatctgggc ctgaacctgc cctgggttgc 6125

ttctgtcctc ag tg gcc aat atc ttt gcc gtg gct gcg ttc cag gtg gag 6175  
Val Ala Asn Ile Phe Ala Val Ala Ala Phe Gln Val Glu  
140 145

aag cgc ctg gcc gtg gtaagcagtg ccctcacgcc ctcccctgac ttgcctcaag 6230  
Lys Arg Leu Ala Val  
150

gtccttacca gtcgggctta gggcgggcca ccagctggtc ccactgtgct tcagggtttt 6290

gggcctttcg tggccttctt gagaggggct gcacctcagg cctggtggct cttcctcagg 6350

gaggtcctct gaccaggag gggggtcctt ggctgacgct ctgctccac cccag gga 6408  
Gly

gct ctg acg gag cag gcg ggg ctg ctg ctg cac ggg gtc aac ctg gcc 6456  
Ala Leu Thr Glu Gln Ala Gly Leu Leu Leu His Gly Val Asn Leu Ala  
155 160 165 170

acc att ctc tgc ttc cca gcg gcc gtg gcc ttt ctc ctc gag tct atc 6504  
Thr Ile Leu Cys Phe Pro Ala Ala Val Ala Phe Leu Leu Glu Ser Ile  
175 180 185

act cca g gtgggcccc cccccgccc cgccccgcc cagctgtct cggccacggg 6561  
Thr Pro

cagcgggggg ggcgtggcct gagcttgccct ctcccacag tg ggc tcc gtg ctg 6614  
Val Gly Ser Val Leu  
190

gcc ctg atg gtc tac acc atc ctc ttc ctc aag ctg ttc tcc tac cgg 6662  
Ala Leu Met Val Tyr Thr Ile Leu Phe Leu Lys Leu Phe Ser Tyr Arg  
195 200 205

gac gtc aac ctc tgg tgc cga gag cgc agg gct ggg gcc aag gcc aag 6710  
Asp Val Asn Leu Trp Cys Arg Glu Arg Arg Ala Gly Ala Lys Ala Lys  
210 215 220 225

gct g gtgagggctg cctcgggctg gggccactgg gctgccactt gcctcgggac 6764  
Ala

cggcaggggc tcggctcacc cccgaccgc cccctgccgc ttgctcgtag ct ttg 6819  
Ala Leu

FIGURE 2b continued

22/34

gca ggt aag aag gcc aac ggg gga gct gcc cag cgc acc gtg agc tac Ala Gly Lys Lys Ala Asn Gly Gly Ala Ala Gln Arg Thr Val Ser Tyr 230 235 240	6867
ccc gac aac ctg acc tac cgc g gtgaggatcc tgccgggggc tggggggact Pro Asp Asn Leu Thr Tyr Arg 245 250	6919
gcccggcggc ctggcctgct agccccgccc tcccttcag at ctc tac tac ttc Asp Leu Tyr Tyr Phe 255	6973
ctc ttc gcc ccc acc ctg tgc tac gag ctc aac ttc ccc cgc tcc ccc Leu Phe Ala Pro Thr Leu Cys Tyr Glu Leu Asn Phe Pro Arg Ser Pro 260 265 270	7021
cgc atc cga aag cgc ttc ctg ctg cgg cga ctc ctg gag atg Arg Ile Arg Lys Arg Phe Leu Leu Arg Arg Leu Leu Glu Met 275 280 285	7063
gtgaggcggg gcctcgcggg ccagggtggg cgggcctgcc ggcacccggc accggggctc agctcactgt ccgcttgctt ccttccccag ctg ttc ctc acc cag ctc cag gtg Leu Phe Leu Thr Gln Leu Gln Val 290	7123 7177
ggg ctg atc cag cag gtacgtgcc gggggggggg gggggggggg gggggggact Gly Leu Ile Gln Gln 295	7232
ctggggccgt tggggagctg actctgcgct ttttgag tgg atg gtc ccg gcc atc Trp Met Val Pro Ala Ile 300 305	7288
cag aac tcc atg aag ccc ttc aag gtgagcaggc aggcctggca ggggtgggtc Gln Asn Ser Met Lys Pro Phe Lys 310	7342
cggggtcagg gctgaggag ccagctgtgc cctgtgccca cag gac atg gac tac Asp Met Asp Tyr 315	7397
tcc cgc atc gtg gag cgc ctc ctg aag ctg gcg gtgagtgacc tgctgggtg Ser Arg Ile Val Glu Arg Leu Leu Lys Leu Ala 320 325	7450
ggacgcgtgg gggcggtgg ggctgttctg gcacctggca cccactcccc acag gtc Val	7507
ccc aac cac ctc atc tgg ctc atc ttc ttc tac tgg ctc ttc cac tcc Pro Asn His Leu Ile Trp Leu Ile Phe Phe Tyr Trp Leu Phe His Ser 330 335 340 345	7555
tgc ctg aac gcc gtg gct gag ctc atg cag ttt gga gac cgc gag ttc Cys Leu Asn Ala Val Ala Glu Leu Met Gln Phe Gly Asp Arg Glu Phe 350 355 360	7603

FIGURE 2b continued



## 23/34

tac cgg gac tgg tg gtgggtggcc ttgccggggc gggggtggtg ggggcccccg 7657  
 Tyr Arg Asp Trp Trp  
 365

ctggggctgg ggccggagcc cctgcccact ctgccccgcc cccgcag g aac tcc gag 7714  
 Asn Ser Glu

tcc atc acc tac ttc tgg cag aac tgg aac atc cct gtt cac aag tgg 7762  
 Ser Ile Thr Tyr Phe Trp Gln Asn Trp Asn Ile Pro Val His Lys Trp  
 370 375 380 385

tgc atc ag gtgggtgtgc gcctgggggc gggggggttg ggggtgggac 7810  
 Cys Ile Arg

ggggtcgcgt ggccccggcg cccagccac tgccgcctcc cccgcag a cac ttc tac 7867  
 His Phe Tyr  
 390

aag ccc atg ctc cgg cgg ggc agc agc aag tgg gca gcc agg acg gca 7915  
 Lys Pro Met Leu Arg Arg Gly Ser Ser Lys Trp Ala Ala Arg Thr Ala  
 395 400 405

gtg ttt ctg gcc tcc gcc ttc ttc cac gag gtcagtgcac tgagggcgcg 7965  
 Val Phe Leu Ala Ser Ala Phe Phe His Glu  
 410 415

ccctgcccct ggtgggggtg ggggtggggg tgggggctcg ctgacgcccc tctcccctca 8025

g tac ctg gtg agc atc ccc ctg cgc atg ttc cgc ctc tgg gcc ttc acc 8074  
 Tyr Leu Val Ser Ile Pro Leu Arg Met Phe Arg Leu Trp Ala Phe Thr  
 420 425 430

ggc atg atg gcg cag gtgagcagcc ctggaccccc gctccgcccc gccccgcgag 8129  
 Gly Met Met Ala Gln  
 435

cgcagaggct cactcccgtc ctgtgtcccc ag atc ccg ctg gcc tgg ata gtg 8182  
 Ile Pro Leu Ala Trp Ile Val  
 440 445

ggc cgc ttc ttc cgc ggc aac tac ggc aac gcg gcc gtg tgg ctg tca 8230  
 Gly Arg Phe Phe Arg Gly Asn Tyr Gly Asn Ala Ala Val Trp Leu Ser  
 450 455 460

ctc atc atc ggg cag ccg gtg gcc gtc ctg atg tac gtc cac gac tac 8278  
 Leu Ile Ile Gly Gln Pro Val Ala Val Leu Met Tyr Val His Asp Tyr  
 465 470 475

tac gtg ctc aac cgt gag gcg ccg gca gcc ggc acc tgagcgccctc 8324  
 Tyr Val Leu Asn Arg Glu Ala Pro Ala Ala Gly Thr  
 480 485

caggctggcc cctcgtggg tgggtgactg ctttgccgcg ctgcctgcgg ctggactaga 8384

gcctgccccca acctggggcg agcaggagga ggcctggctg gtggaagctg cctcctggcc 8444

tccaccaggc ctctgcctga agggcttcct cctgccaggg gagagcaggc ccgacgcagt 8504

tctggcccct gggaggtgcc catgctctg aaacctaca gatctcgccc aagggtctga 8564

FIGURE 2b continued

24/34

atgtgtcaat aaagtgtgtg gcacagttag ctccctcagc ctccagggca cagggtgtgc 8624  
 aggagggggc ggccctccca cgtggggcca tgctgtggga aggagggccc agcgctgtga 8684  
 gaggagtgtg ggctgtgtg accctccctg cctcacaggg ctctgtgtgc agacgtcttg 8744  
 cctgtcaagg tggagactcc atgctccaag gccccctgtg cctgaggtct gcacacaagt 8804  
 ggattcaact tgggtcaggc cagaggctaa ggtgtggaag aagggtgaga atcaggctga 8864  
 cttgaacggc agcaaagact ccaaggcaag gctgcagaag tctcaaaagc tatgcgcaca 8924  
 gtccctgtct ggggtgtctc cctgggtgtg gctctgggtc gcttgacaa agcaggtgtc 8984  
 ctgggtcagc cctcaccgag ggccttcctt gggggcagag gttggcctga tgccaggggc 9044  
 tccccgtttt tccaggccct cagcaggtag ttgggtgtgg ccctcatgat acctgtgtcc 9104  
 cagagctccg ccaactaaaa agcttggcag tgaggcaagg gcaaccccg gctgttcccc 9164  
 cctctactgg ctctgccgcc tgggttgga accctgaggc tgtgccaggc aggtgtacct 9224  
 tgacagccag ccatggccca gtaagatggg tgcccaggtt ggtacctggg cagcggacct 9284  
 agctgtgtctg ccccgcccc aaccagaagc cgctctagcc catggtgtgtc gtctgggcga 9344  
 gacaggtgtg ttggctaggc actgtttgtt ctacagcagg tgtaggcagc gtctccctga 9404  
 cccctgcctt ctaggaagcc accaccctgn gccctactca tcagcaagga cagcgagcag 9464  
 ggctgagctg ggggtgcgtg ggctgtctac gnccccgcac ctctcatcaca tgcacctctg 9524  
 caccctctgc tgctgactc aagagtgggg ggggggttcc tgtgttctt tcattccaga 9584  
 cccacgtgtc tgacctagt caccacctg gtccgtagt gctgacctg ccacagggct 9644  
 cctgtgggccc cagctgctc ccgcctgtt cccttcataa agaactcttg agcacatgca 9704  
 gccagggga gccagaggc tccagtgtgc tgtgtccatc tgcctccctt cagcccttc 9764  
 cgagacactg cgcatcatgc cccctccac cccacccac actggcagga ggaacacaca 9824  
 gggagaccac acacagagct cgttgtttat aaatctctgc ctggctcatc ggtctgtttg 9884  
 tccatgtata tatctgcata tctctatgga aggggaaagg gggactcgtg taaaaatcca 9944  
 aaatacaatt ctatgaacac ctgcatcctg gtcagtctga gttgtggcgt gaagcccagg 10004  
 tgagctgtgg ctcacaggc taggcctctg gtgtgtggcg ggggacctc cccacccct 10064  
 ctccccctc ccgccagcca ggggaccagg ctctggaca ccaggcctgc ccaaggcctg 10124  
 ctctcctctt ggggttctc cgagacagt gggctcttgg ctttggggg ttctgagccc 10184  
 gtcagcagg agatgtgtg gtcattctct atatttcgtc tccctcgag aagtaggagc 10244  
 cctccccag ctgaagagc accggcaggt cgctgtctcc cagctccag gagcccggt 10304  
 ccaggagcag caggggtgtg gcgtgtagt gcaccagctg ctccctagg ggtgcgactg 10364

FIGURE 2b continued

25/34

ggtcaggggtg ccggtggggc cggggggcgg ggtggggnnn nnnnnnnnnn nnncccccc 10424  
ccccggcccc agccaccccg cctacgcacg ctggccaggc tgctgtccag gtcgggcagg 10484  
ctcatgtcgg gcaccgtaac cgaggggctg aacagctgca gggaagaggg gcgggtcaga 10544  
ctgccctgga agcggggcgg gctgggcgcg gggcgggcag ggtagactc acatccagca 10604  
gggtgctggt gtccacgctg aagccatggc ttgtcagcat ggtctgcagg ttgtccagg 10664  
tggagtccat ggcgtccaag tggtcgctga gctcggctct ggccagagga aggggagcag 10724  
gtgacgtggc atccaggcac cccaggtgc agtcctgccg gcccttaagc ccagctgcca 10784  
gcctgccctg cgcaggctca cgggaggccc tgaggtggg ccctgggcct gggacacttg 10844  
gccaccggtg gtggctcagg ctccctcacc accgtgaggc ccgtaaggc agcacgtcag 10904  
gctgggcccg ctaagacatc agggggctga gttcagggtc cagaagggcc cggggcccca 10964  
gcttctcagg ttcagcaacg cctgcccttc aactggagg acagcaggag ggtggagctg 11024  
tgctcgcca ggctgggcca gcctacccca gcatgcgcca ggatagatta ggcctgcccc 11084  
ctcgaggagg ccaggcagta cctcggggtg ctgcctcagg cccagagggt ggggtgggag 11144  
cctgttcaca ggctgggag ggggtctgctc agaggctggg gaggggcccc agctctggcg 11204  
gagccctcct ccctccccag tgggacagcg ctaaccctgg ctggactcgg ccatgcagag 11264  
aggaggaggg gcaggggaag aggcgggcga cccagacat ctttgagtg cgagccaaac 11324  
tgcaagatag aaagacaaga gccccctct gcgggtggg cccagcgagt ccaagcccc 11384  
caccctgttc acccacctct ggggtggtg gacagacaga ccaggcgcc cggggagggc 11444  
ggctcagggg gcagggtgc ccgccacgtg cctggttccc acggtgcagg ggctgcggga 11504  
ccccctggct ggcattgtct cctccccagc tccatgtctt gggccaggtc ggccagtttc 11564  
cccagcaaa tcccatgccg agggcctgag ggccatgct gtccagcccg gcaactgtct 11624  
tgtcggaggc ccctgcctt ccagagaccg aaggcgccct ggaaggcact cacttgtota 11684  
ggcaggcgac gctgaggcac ttctcgggag ccgaggcggg gagggtcgac ggatagcggc 11744  
cccgagtgat ccgatagaag cttcgta 11771

FIGURE 2b continued

26/34




FIGURE 3

27/34

&lt;221&gt; VARIANT

&lt;222&gt; (232)..(232)

<223> an amino acid substitution (K -> A) caused by a polymorphism at bases 7224-7225 of the genomic sequence (measured from the adenine residue of the translation start codon). Lysine (K) corresponds to the Q allele, alanine (A) corresponds to the q allele.

&lt;400&gt; 5

Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Arg Thr Gly  
1 5 10 15

Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu  
20 25 30

Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr  
35 40 45

Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asp Leu Arg  
50 55 60

Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser Asp Ser Gly Phe Ser  
65 70 75 80

Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val Met Leu Ile Leu Ser  
85 90 95

Asn Ala Arg Leu Phe Leu Glu Asn Leu Ile Lys Tyr Gly Ile Leu Val  
100 105 110

Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro Tyr Ser Trp  
115 120 125

Pro Ala Leu Cys Leu Val Ile Val Ala Asn Ile Phe Ala Val Ala Ala  
130 135 140

Phe Gln Val Glu Lys Arg Leu Ala Val Gly Ala Leu Thr Glu Gln Ala  
145 150 155 160

Gly Leu Leu Leu His Gly Val Asn Leu Ala Thr Ile Leu Cys Phe Pro  
165 170 175

FIGURE 4a

28/34

Ala Ala Val Ala Phe Leu Leu Glu Ser Ile Thr Pro Val Gly Ser Val  
180 185 190

Leu Ala Leu Met Val Tyr Thr Ile Leu Phe Leu Lys Leu Phe Ser Tyr  
195 200 205

Arg Asp Val Asn Leu Trp Cys Arg Glu Arg Arg Ala Gly Ala Lys Ala  
210 215 220

Lys Ala Ala Leu Ala Gly Lys Lys Ala Asn Gly Gly Ala Ala Gln Arg  
225 230 235 240

Thr Val Ser Tyr Pro Asp Asn Leu Thr Tyr Arg Asp Leu Tyr Tyr Phe  
245 250 255

Leu Phe Ala Pro Thr Leu Cys Tyr Glu Leu Asn Phe Pro Arg Ser Pro  
260 265 270

Arg Ile Arg Lys Arg Phe Leu Leu Arg Arg Leu Leu Glu Met Leu Phe  
275 280 285

Leu Thr Gln Leu Gln Val Gly Leu Ile Gln Gln Trp Met Val Pro Ala  
290 295 300

Ile Gln Asn Ser Met Lys Pro Phe Lys Asp Met Asp Tyr Ser Arg Ile  
305 310 315 320

Val Glu Arg Leu Leu Lys Leu Ala Val Pro Asn His Leu Ile Trp Leu  
325 330 335

Ile Phe Phe Tyr Trp Leu Phe His Ser Cys Leu Asn Ala Val Ala Glu  
340 345 350

Leu Met Gln Phe Gly Asp Arg Glu Phe Tyr Arg Asp Trp Trp Asn Ser  
355 360 365

Glu Ser Ile Thr Tyr Phe Trp Gln Asn Trp Asn Ile Pro Val His Lys  
370 375 380

Trp Cys Ile Arg His Phe Tyr Lys Pro Met Leu Arg Arg Gly Ser Ser  
385 390 395 400

Lys Trp Ala Ala Arg Thr Ala Val Phe Leu Ala Ser Ala Phe Phe His  
405 410 415

FIGURE 4a continued

29/34

Glu Tyr Leu Val Ser Ile Pro Leu Arg Met Phe Arg Leu Trp Ala Phe  
420 425 430

Thr Gly Met Met Ala Gln Ile Pro Leu Ala Trp Ile Val Gly Arg Phe  
435 440 445

Phe Arg Gly Asn Tyr Gly Asn Ala Ala Val Trp Leu Ser Leu Ile Ile  
450 455 460

Gly Gln Pro Val Ala Val Leu Met Tyr Val His Asp Tyr Tyr Val Leu  
465 470 475 480

Asn Arg Glu Ala Pro Ala Ala Gly Thr  
485

FIGURE 4a continued

30/34

&lt;400&gt; 6

Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Arg Thr Gly  
 1 5 10 15

Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu  
 20 25 30

Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr  
 35 40 45

Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asp Leu Arg  
 50 55 60

Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser Asp Ser Gly Phe Ser  
 65 70 75 80

Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val Met Leu Ile Leu Ser  
 85 90 95

Asn Ala Arg Leu Phe Leu Glu Asn Leu Ile Lys Tyr Gly Ile Leu Val  
 100 105 110

Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro Tyr Ser Trp  
 115 120 125

Pro Ala Leu Cys Leu Val Ile Val Ala Asn Ile Phe Ala Val Ala Ala  
 130 135 140

Phe Gln Val Glu Lys Arg Leu Ala Val Gly Ala Leu Thr Glu Gln Ala  
 145 150 155 160

Gly Leu Leu Leu His Gly Val Asn Leu Ala Thr Ile Leu Cys Phe Pro  
 165 170 175

Ala Ala Val Ala Phe Leu Leu Glu Ser Ile Thr Pro Val Gly Ser Val  
 180 185 190

Leu Ala Leu Met Val Tyr Thr Ile Leu Phe Leu Lys Leu Phe Ser Tyr  
 195 200 205

Arg Asp Val Asn Leu Trp Cys Arg Glu Arg Arg Ala Gly Ala Lys Ala  
 210 215 220

Lys Ala Ala Leu Ala Asp Leu Tyr Tyr Phe Leu Phe Ala Pro Thr Leu  
 225 230 235 240

FIGURE 4b



## 31/34

Cys Tyr Glu Leu Asn Phe Pro Arg Ser Pro Arg Ile Arg Lys Arg Phe  
245 250 255

Leu Leu Arg Arg Leu Leu Glu Met Leu Phe Leu Thr Gln Leu Gln Val  
260 265 270

Gly Leu Ile Gln Gln Trp Met Val Pro Ala Ile Gln Asn Ser Met Lys  
275 280 285

Pro Phe Lys Asp Met Asp Tyr Ser Arg Ile Val Glu Arg Leu Leu Lys  
290 295 300

Leu Ala Val Pro Asn His Leu Ile Trp Leu Ile Phe Phe Tyr Trp Leu  
305 310 315 320

Phe His Ser Cys Leu Asn Ala Val Ala Glu Leu Met Gln Phe Gly Asp  
325 330 335

Arg Glu Phe Tyr Arg Asp Trp Trp Asn Ser Glu Ser Ile Thr Tyr Phe  
340 345 350

Trp Gln Asn Trp Asn Ile Pro Val His Lys Trp Cys Ile Arg His Phe  
355 360 365

Tyr Lys Pro Met Leu Arg Arg Gly Ser Ser Lys Trp Ala Ala Arg Thr  
370 375 380

Ala Val Phe Leu Ala Ser Ala Phe Phe His Glu Tyr Leu Val Ser Ile  
385 390 395 400

Pro Leu Arg Met Phe Arg Leu Trp Ala Phe Thr Gly Met Met Ala Gln  
405 410 415

Ile Pro Leu Ala Trp Ile Val Gly Arg Phe Phe Arg Gly Asn Tyr Gly  
420 425 430

Asn Ala Ala Val Trp Leu Ser Leu Ile Ile Gly Gln Pro Val Ala Val  
435 440 445

Leu Met Tyr Val His Asp Tyr Tyr Val Leu Asn Arg Glu Ala Pro Ala  
450 455 460

Ala Gly Thr  
465

FIGURE 4b continued

32/34

K232A



*Bos taurus*: LALMVYTIILFLKFSYRDVNLWCRRERAGAKAKAALAGKKANGGAAQRTVSPDNLTYRDLYYFLFAPTLCY  
*Bison bison*: LALMVYTIILFLKFSYRDVNLWCRRERAGAKAKAALAGKKANGGAAQRTVSPDNLTYRDLYYFLFAPTLCY  
*Ovis aries*: LALMVYTIILFLKFSYRDVNLWCRRERAGAKAKAALAGKKANGGAAQRTVSPDNLTYRDLYYFLFAPTLCY  
*Sus scrofa*: LALMVYAIILFLKFSYRDVNLWCRRERATAKAASAGKKANGGAAQHSVSPDNLTYRDLYYFLFAPTLCY  
*Homo sapiens*: LALMAHTIILFLKFSYRDVNSWC---RR---ARAKASAGKKASAAAPHVSPDNLTYRDLYYFLFAPTLCY  
*Cercopithecus aethiops*: LALMVHTIILFLKFSYRDVNLWC---RR---ARAKASAGKRASAAAPHVSPDNLTYRDLYYFLFAPTLCY  
*M. musculus domesticus*: FALASYSIMFLKLYSYRDVNLWCRR--VKAKAVSTGKKVSGAAQQAVSPDNLTYRDLYYFIFAPTLCY  
*Rattus norvegicus*: FALASYSIILFLKSSYRDVNLWCRR--VKAKAVSAGKKVSGAAQQAVSPDNLTYRDLYYFIFAPTLCY

FIGURE 5




FIGURE 6

34/34




FIGURE 7

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ01/00245

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>																						
Int. Cl. <sup>7</sup> : C12Q 1/68, C12N 15/12, A01K 67/027, C07K 16/40																						
According to International Patent Classification (IPC) or to both national classification and IPC																						
<b>B. FIELDS SEARCHED</b>																						
Minimum documentation searched (classification system followed by classification symbols)																						
WPIDS, CA: SEE BELOW																						
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched																						
AGRICOLA, BIOSIS, DNA DATABASES (GENBANK, EMBL): SEE BELOW																						
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)																						
WPIDS, GENBANK, EMBL, BIOSIS, CA, MEDLINE, AGRICOLA: diacylglycerol acyltransferase, dgat1, bovine, cow, cattle, milk																						
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>																						
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																				
X	EMBL Accession No AF059202 15 October 1998 Oelkers P et al <i>Homo sapiens</i> ACAT related gene product 1 mRNA, complete cds	27-33, 35-37																				
X	EMBL Accession No AF236018 12 August 2000 Joyce CW et al <i>Cercopithecus aethiops</i> diacyl-glycerol acyltransferase mRNA, complete cds	27-33, 35-37																				
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input type="checkbox"/> See patent family annex																						
<p>* Special categories of cited documents:</p> <table border="0"> <tr> <td>"A"</td> <td>document defining the general state of the art which is not considered to be of particular relevance</td> <td>"T"</td> <td>later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"E"</td> <td>earlier application or patent but published on or after the international filing date</td> <td>"X"</td> <td>document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"I"</td> <td>document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"Y"</td> <td>document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"O"</td> <td>document referring to an oral disclosure, use, exhibition or other means</td> <td>"&amp;"</td> <td>document member of the same patent family</td> </tr> <tr> <td>"P"</td> <td>document published prior to the international filing date but later than the priority date claimed</td> <td></td> <td></td> </tr> </table>			"A"	document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"E"	earlier application or patent but published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"I"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family	"P"	document published prior to the international filing date but later than the priority date claimed		
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																			
"E"	earlier application or patent but published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																			
"I"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																			
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family																			
"P"	document published prior to the international filing date but later than the priority date claimed																					
Date of the actual completion of the international search		Date of mailing of the international search report																				
5 February 2002		13 FEB 2002																				
Name and mailing address of the ISA/AU		Authorized officer																				
AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustalia.gov.au Facsimile No. (02) 6285 3929		<b>TERRY MOORE</b> Telephone No : (02) 6283 2632																				

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ01/00245

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBL Accession No AF078752 11 November 1998 Cases S et al <i>Mus musculus</i> diacylglycerol acyltransferase (Dgat) mRNA, complete cds	27-33, 35-37
X	Smith CJ et al "Obesity resistance and multiple mechanisms of triglyceride synthesis in mice lacking Dgat" (2000) Nature Genetics 25(1), pages 87-90 See the whole document.	44-46

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ01/00245

**Box I** Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos :  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos : 1-50 in part and claims 7 and 14-22 in full.  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
See the explanation in the supplemental box.
3. ☐ Claims Nos :  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

**Box II** Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ01/00245

**Supplemental Box**

(To be used when the space in any of Boxes I to VIII is not sufficient)

**Continuation of Box No: Box I(2)**

Claims 1-50 have not been searched in full because the applicant, although invited to furnish the sequence listing in computer readable form, failed to do so. As such the claims were not searched with regard to the sequences disclosed in the specification. The claims have simply been searched using prior art human and mouse DGAT1 gene sequences located in the EMBL and GENBANK databases.

Claims 7 and 14-22 have not been searched in full because the scope of the claims is unclear. The claims include within their scope bovines that are not the products of the invention and whose pre-existing genotype is simply further characterised using the DGAT1 gene. It is impossible to determine which bovines would fall within the scope of these claims and as such a meaningful search cannot be conducted.